


HARFORD COUNTY

RESULTS OF THE 1994-1997 MARYLAND BIOLOGICAL STREAM SURVEY: COUNTY ASSESSMENTS



**CHESAPEAKE BAY AND
WATERSHED PROGRAMS**
MONITORING AND
NON-TIDAL ASSESSMENT
CBWP-MANTA-EA-01-21





Parris N. Glendening
Governor

Kathleen Kennedy Townsend
Lt. Governor

A message to Maryland's citizens

The Maryland Department of Natural Resources (DNR) seeks to preserve, protect and enhance the living resources of the state. Working in partnership with the citizens of Maryland, this worthwhile goal will become a reality. This publication provides information that will increase your understanding of how DNR strives to reach that goal through its many diverse programs.

J. Charles Fox
Secretary

Karen M. White
Deputy Secretary



Maryland Department of Natural Resources
Tawes State Office Building
580 Taylor Avenue
Annapolis, Maryland 21401

Toll free in Maryland: 1-(877) 620 8DNR x8611
Out of state call: 410-260-8611
www.dnr.state.md.us

The facilities and services of the Maryland Department of Natural Resources are available to all without regard to race, color, religion, sex, sexual orientation, age, national origin, physical or mental disability.

This document is available in alternative format upon request from a qualified individual with a disability.

Publication date: December 2001

© MD DNR 2001



PRINTED ON RECYCLED PAPER

HARFORD COUNTY

Results of the 1994-1997 Maryland Biological Stream Survey: County-Level Assessments

**Christopher J. Millard
Paul F. Kazyak
Anthony P. Prochaska**

December 2001

**Maryland Department of Natural Resources
Resource Assessment Service
Monitoring and Non-Tidal Assessment Division
580 Taylor Avenue
Annapolis, MD 21401**

**THIS PAGE INTENTIONALLY
LEFT BLANK**

FOREWORD

This report is based on results of the Maryland Biological Stream Survey (MBSS), a program funded primarily by the Power Plant Research Program and administered by the Maryland Department of Natural Resources (MDNR). Field data for the MBSS were collected by the Maryland Department of Natural Resources. Analyses of water chemistry samples were conducted by the University of Maryland's Appalachian Laboratory. Much of the initial data analysis was conducted by Versar, Inc. for MDNR's Power Plant Assessment Division.

This report helps fulfill two outcomes in MDNR's Strategic Plan: 1) A Vital and Life Sustaining Chesapeake Bay and Its Tributaries, and 2) Sustainable Populations of Living Resources and Healthy Ecosystems.

ACKNOWLEDGMENTS

The 1994-1997 Maryland Biological Stream Survey has been a cooperative effort among several agencies, consultants and academic institutions. We wish to thank Nancy Roth and Ginny Mercurio from Versar in helping to compile some of the data used in this report. Versar also designed the sampling program, obtained landowners' permissions, and helped manage the data. We are also grateful to the many individuals from Maryland Department of Natural Resources, the University of Maryland's Appalachian Laboratory (AL), and the University of Maryland's Wye Research and Education Center (WREC) who comprised the field crews and did a great job collecting the data. MDNR staff also digitized watersheds and calculated land use data, provided quality assurance, and conducted field crew training. Nancy Roth and her colleagues at Versar developed the fish Index of Biotic Integrity, and Dr. Sam Stribling and his staff at Tetra Tech, Inc. developed the benthic Index of Biotic Integrity. Dr. Ray Morgan of AL and Mr. Lenwood Hall of the WREC supervised additional field crews and developed the Physical Habitat Index, and Dr. Keith Eshleman of AL assisted with analyses of data on acidified streams. Drs. Wayne Starnes and Bob Reynolds of the Smithsonian Institution (reptiles and amphibians), Dr. Rich Raesly of Frostburg State University (fish), Rita Villella of the U.S. Geological Survey Leetown Science Center (mussels), and Michael Naylor of MDNR (aquatic vegetation) provided taxonomic verifications of voucher specimens. The success of the project resulted from the strong efforts of all these groups. Special thanks go to Ron Klauda for his editorial support and Brenda Morgan for her assistance in formatting, editing, and organizing the report.

**THIS PAGE INTENTIONALLY
LEFT BLANK**

Table of Contents

FOREWORD	i
ACKNOWLEDGEMENTS	i
INTRODUCTION	1
METHODS	3
MARYLAND BIOLOGICAL STREAM SURVEY DATA	9
COUNTY SUMMARY	10
LIST OF TABLES	
Table 1. Site information and land use data collected at Maryland Biological Stream Survey sites in Harford County, 1994-1997.	11
Table 2. Percent occurrence of fish species collected at Maryland Biological Stream Survey sites in Harford County, 1994-1997.	15
Table 3. Tolerance Value, Functional Feeding Group, Habit, and Percent Occurrence of benthic macroinvertebrate taxa collected at Maryland Biological Stream Survey sites in Harford County, 1994-1997.	17
Table 4. Percent occurrence of reptile and amphibian species collected at Maryland Biological Stream Survey sites in Harford County, 1994-1997.	22
Table 5. Physical habitat data for Maryland Biological Stream Survey sites in Harford County, 1994-1997.	23
Table 6. Fish Index of Biotic Integrity (F-IBI), Benthic Macroinvertebrate Index of Biotic Integrity (B-IBI), Family-Level Benthic Macroinvertebrate Index of Biotic Integrity (Fam. IBI), and Physical Habitat Index (PHI) scores at Maryland Biological Stream Survey sites in Harford County, 1994-1997.	26
Table 7. Water chemistry data collected at Maryland Biological Stream Survey sites in Harford County, 1994-1997.	28
LIST OF FIGURES	
Figure 1. Land use in Harford County.	13
Figure 2. Location of Maryland Biological Stream Survey sites in Harford County, 1994-1997.	14
Figure 3. Stream ecological conditions based on the Fish Index of Biotic Integrity (F-IBI) at Maryland Biological Stream Survey sites in Harford County.	16
Figure 4. Stream ecological conditions based on the Benthic Macroinvertebrate Index of Biotic Integrity (B-IBI) at Maryland Biological Stream Survey sites in Harford County.	21
Figure 5. Stream ecological conditions based on the Physical Habitat Index (PHI) at Maryland Biological Stream Survey sites in Harford County.	25
Figure 6. Nitrate-nitrogen concentrations (mg/L) at Maryland Biological Stream Survey sites in Harford County.	30
LITERATURE CITED	31
APPENDIX A	
Summary of the types of data collected at Maryland Biological Stream Survey sites in Harford County, 1994-1997.	A-1
APPENDIX B	
Physical habitat condition measured by the Maryland Biological Stream Survey, 1994-1997.	B-1

**THIS PAGE INTENTIONALLY
LEFT BLANK**

INTRODUCTION

This report presents county-level data from the 1994–1997 Maryland Biological Stream Survey (MBSS or the Survey). Previous reports have documented interim results from the 1995 (Roth et al. 1997) and 1996 (Roth et al. 1998a) sample years. In addition, a comprehensive final report was produced to assess the “state of the streams” throughout the state (Roth et al. 1999). All previous MBSS reports have presented information by individual drainage basins. Because there is a recognized need for stream health information at the county level, a series of reports were prepared; this report is part of that series. This introductory section recounts the origin of the Survey and describes its components.

Origin of the MBSS

More than 10 years ago, the Maryland Department of Natural Resources (MDNR) recognized that atmospheric deposition was one of the most important environmental problems resulting from the generation of electric power. To determine the extent of acidification of Maryland streams resulting from acidic deposition, MDNR conducted the Maryland Synoptic Stream Chemistry Survey (MSSCS) in 1987. The MSSCS estimated the number and extent of streams at that time affected by or sensitive to acidification statewide and demonstrated the potential for adverse effects on biota from acidification. However, little direct information was available on the biological responses of Maryland streams to water chemistry conditions. Data that were available could not be used (because of methodological differences and spatial coverage limitations) to compare conditions across regions or watersheds (Tornatore et al. 1992). Neither was it possible to assess the interactions between acidic deposition and other anthropogenic and natural influences (CBRM 1989). For these reasons, in 1993, MDNR created the MBSS to provide comprehensive information on the status of biological resources in Maryland streams and how they are affected by acidic deposition and other cumulative effects of anthropogenic stresses.

Description of the MBSS

The MBSS is intended to help environmental decision-

makers protect and restore the natural resources of Maryland. The primary objectives of the MBSS are:

- to assess the current status of biological resources in Maryland’s non-tidal streams;
- to quantify the extent to which acidic deposition has affected or may be affecting biological resources in the state;
- to examine which other water chemistry, physical habitat, and land use factors are important in explaining the current status of biological resources in streams;
- to compile the first statewide inventory of stream biota;
- to establish a benchmark for long-term monitoring of trends in these biological resources; and
- to target future local-scale assessments and mitigation measures needed to restore degraded biological resources.

In creating the Survey, MDNR implemented a probability-based sampling design as a cost-effective way to characterize statewide stream resources. By randomly selecting sites, the Survey can make quantitative inferences about the characteristics of all 9,258 miles of first-to-third-order, non-tidal streams in Maryland (based on stream length on a 1:250,000-scale base map). MDNR recognized that the utility of these estimates depended on accurately measuring appropriate attributes of streams. The Survey focuses on biology for two reasons: (1) organisms themselves have direct societal value and (2) biological communities integrate stresses over time and are a valuable and cost-effective means of assessing ecological integrity (i.e., the capacity of a resource to sustain its inherent potential).

Fish are an important component of stream integrity and one that also contributes to substantial recreational values. For these reasons, fish communities are a primary focus of the Survey. The Survey collects quantitative data for the calculation of population estimates for individual fish species (both game and nongame). These data can also be used to evaluate

fish community composition, individual fish health, and the geographic distribution of commercially important, rare, or non-indigenous fish species. Benthic (bottom-dwelling) macroinvertebrates are another essential component of streams and they constitute the second principal focus of the Survey. The Survey uses rapid bioassessment procedures for collecting benthic macroinvertebrates; these semi-quantitative methods permit comparisons of relative abundance and community composition, and have proven to be an effective way of assessing biological integrity in streams (Hilsenhoff 1987, Lenat 1988, Plafkin et al. 1989, Kerans and Karr 1994, Resh 1995). The Survey also records the presence of reptiles and amphibians (herpetofauna), freshwater mussels, and aquatic plants (both submerged aquatic vegetation (SAV) and emergent macrophytes). The Survey has established rigorous protocols (Kazyak 1996) for each of these sampling components, as well as training and auditing procedures to assure that data quality objectives are met.

Although the MBSS sampling design and protocols provide exceptional information for characterizing the stream resources in Maryland, designation of degraded areas and identification of likely stresses requires additional activities. Assessing the condition of biological resources (whether they are degraded or not degraded) requires the development of ecological indicators that permit the comparison of sampled segment results to minimally impacted reference conditions (i.e., the biological community expected in watersheds with little or no human-induced impacts). The Survey has used its growing database of information collected with consistent methods and broad coverage across the state to develop and test indicators of individual biological components (Stribling et al. 1998, Roth et al. 1998b) and physical habitat quality (Hall et al. 1999). Each of these indicators consists of multiple metrics using the general approach developed for the Index of Biotic Integrity (IBI) (Karr et al. 1986, Karr 1991) and the Chesapeake Bay Benthic Restoration Goals (Ranasinghe et al. 1994). The fish and benthic macroinvertebrate IBIs (which combine attributes of both the number and the type of species found) are widely accepted indicators that have been adapted for use in a variety of geographic locations (Miller et al. 1988, Cairns and Pratt 1993, Simon 1999). The Survey is investigating the possibility

of developing additional indicators (e.g., amphibians in small streams with few or no fish) and combining components into a composite indicator of biological integrity.

In addition to developing reference-based indicators, the Survey is applying a variety of analytical methods to the question of which stressors are most closely associated with degraded streams. This involves correlational and multivariate analyses of water chemistry, physical habitat, land use, and biological information (e.g., presence of non-native species). The biological information also provides a valuable opportunity for documenting aquatic biodiversity across the state; the distribution and abundance of species previously designated as rare only by anecdotal evidence can be determined, and unique combinations of species at the ecosystem and landscape levels can be identified. Land use and other landscape-scale metrics will play an important role in identifying the relative contributions of different stressors to the cumulative impact on stream resources. Ultimately, the Survey seeks to provide an integrated assessment of the problems facing Maryland streams that will facilitate interdisciplinary solutions for their restoration. The survey also provides resource managers with the locations of relatively undisturbed streams and watersheds that deserve protection.

METHODS

This section presents the specific study design and procedures used to implement the Maryland Biological Stream Survey. The study area of concern and the sampling design developed to characterize it are presented, along with field and laboratory methods for each component: fish, benthic macroinvertebrates, reptiles and amphibians, physical habitat, and water chemistry. Methods for aquatic vegetation and mussel sampling are presented, but the resulting data are not included in this report. A full description of MBSS methods can be found in Kazyak (1996).

MBSS Study Design

The Survey study area comprises 17 distinct drainage basins across the state. Random sampling was used to allow the estimation of unbiased summary statistics (e.g., means, proportions, and their respective variances) for the entire state, a particular basin, and subpopulations of interest (e.g., streams with pH < 5).

Because it would have been cost prohibitive to visit a sufficient number of sites in all basins in a single year, lattice sampling was used to schedule sampling of all basins over a three-year period, 1995-1997. Lattice sampling, also known as multistratification, is a cost-effective means of allocating effort across time in a large geographic area (Heimbuch 1999, Jessen 1978, Cochran 1977). A table, or lattice, was formed by arranging 17 basins in 17 rows, and the years in 3 columns. Lattice sampling was the method used for selecting cells from this 17x3 table so that all basins would be sampled over a three-year period and all basins would have a non-zero probability of being sampled in a given year. The data presented in this report include those collected at random sampling sites within the 17 principal basins in Maryland, as well as sites from the 1994 demonstration project. Because no estimates were calculated for this report, these data were included to supplement the number of sites.

The sampling frame for the Survey was constructed by overlaying basin boundaries on a map of all blue-line stream reaches in the study area as digitized on a U.S. Geological Survey 1:250,000 scale topographic map. This sample frame was similar to that used by the earlier Maryland Synoptic Stream Chemistry Survey

(MSSCS) conducted in 1987 (Knapp and Saunders 1987, Knapp et al. 1988). The Strahler convention (Strahler 1957) was used for ranking stream reaches by order; first-order reaches, for example, are the most upstream reaches in the branching stream system. Sampling was restricted to non-tidal, third-order and smaller stream reaches, excluding impoundments that were non-wadable or that substantially altered the riverine nature of the reach (Kazyak 1994). Together, these first-through third-order streams comprise about 90% of all stream and river miles in Maryland. Stream reaches were further divided into non-overlapping, 75-meter segments; these segments were the elementary sampling units from which biological, water chemistry, and physical habitat data were collected.

The 1995-1997 MBSS study design was based on stratified random sampling of segments within each basin; each basin was stratified by stream order. Within a stream order, the number of segments sampled per basin is proportional to the number of stream miles in the basin. To achieve the target number of samples per stream order within each basin, a given number of segments were randomly selected from each basin and ranked in order of selection. In all basins, extra segments were selected as a contingency against loss of sampling sites from restricted access to selected streams or from streams that were dry, too deep, or otherwise unsampleable owing to field conditions. In some basins, where only a small number of sites would have been selected using this method, additional random sites were selected to increase sample size. These extra sites (selected at random using the method described above) were used to provide better basinwide estimates; they were not included in the estimates of statewide conditions.

Permissions were obtained to access privately owned land adjacent to or near each stream segment. The procedures for obtaining permissions are described in Chaillou (1995). Because landowner permissions were obtained in a synoptic fashion and some variation in these rates occurred, we obtained more permissions than were needed for the Survey. Only the highest ranking sites were sampled until the target goal for that basin was reached. For the three year study, the success rate for obtaining permission to access stream sampling segments was high. Eighty-eight percent of sites that were targeted for permission were sampled.

Reasons for permission denial varied and generally reflected the preferences of landowners regarding property access, rather than any specific types of land. In rare cases, permission denial may affect the interpretation of Survey estimates, but only where denials occur in streams with characteristics that differ from the general population of streams. In one example of potential bias, several sites with known coal mining activities in the North Branch Potomac basin denied permission to sample, likely under representing the proportion of acid mine drainage streams in the population.

Field and Laboratory Methods

Benthic macroinvertebrate and water quality sampling were conducted in spring, when the benthos are thought to be reliable indicators of environmental stress (Plafkin et al. 1989) and when acid deposition effects are often the most pronounced. Fish, reptiles and amphibians, aquatic vegetation, and mussel sampling, along with physical habitat evaluations, were conducted during the low-flow period in summer. Fish community composition tends to be stable during summer, and low flow is advantageous for electrofishing. Because low-flow conditions in summer may be a primary factor limiting the abundance and distribution of fish populations, habitat assessments were performed during the summer. The sample size in summer is lower than in spring because some streams were dry in summer or were, in rare cases, otherwise unsampleable.

To reduce temporal variability, sampling during spring and summer was conducted within specific, relatively narrow time intervals, referred to as index periods (Janicki et al. 1993). These index periods were defined by degree-day limits for specific parts of the state. This approach provided a synoptic assessment of the current status of stream biota, water quality, and physical habitat in the 17 basins sampled. The spring index period was the time period between approximately March 1 and May 1, with end of the index period determined by degree-day accumulation as specified in Hilsenhoff (1987). In reality, most spring samples (78%) were collected in March, well before degree-day accumulation limits were approached. The summer index period was between June 1 and September 30 (Kazyak 1994).

Data Collection and Measurement

Field sampling followed procedures specified in the MBSS sampling manual (e.g., Kazyak 1996). A summary of the variables measured and the field and laboratory methods used to conduct the sampling follows.

Fish

Fish were sampled during the summer index period using double-pass electrofishing within 75-meter stream segments. Block nets were placed at each end of the segment and direct current backpack electrofishing units were used to sample the entire segment. An attempt was made to thoroughly fish each segment, and consistent effort was applied over the two passes. This sampling approach allowed calculation of several metrics useful in calculating a biological index and produced unbiased estimates of fish species abundance.

In small streams, a single electrofishing unit was used. In larger streams, two to five units were employed to effectively sample the site. Captured fish were identified to species, counted, weighed, and released. Any individuals that could not be identified to species were retained for laboratory confirmation. For each pass, all individuals of each gamefish species (defined as trout, bass, walleye, pike, chain pickerel, and striped bass) were measured for total length and examined for visible external pathologies or anomalies. For nongame species, up to 100 fish of each species (from both passes) were examined for visible external pathologies or anomalies. For each pass, all non-game species were weighed together for an aggregate biomass measurement; gamefish were also weighed in aggregate to the nearest 10 g.

Electrofishing was also conducted at supplemental, non-randomly selected sites during the summer index period. The presence of each species of fish was recorded for these segments to provide additional qualitative information on statewide fish distributions. Sampling effort at most qualitative sites was based on doubling the elapsed time since the last species was recorded or a minimum of 600 seconds of electrofishing effort.

After processing the fish collected in the field, voucher

specimens were retained for each species not previously collected in the drainage basin. In addition, all individuals which could not be positively identified in the field were retained. The remaining fish were released. All voucher specimens and fish retained for positive identification in the laboratory were examined and verified by the MBSS Quality Assurance Officer or ichthyologists at Frostburg State University, Frostburg, Maryland or the Smithsonian Institution, Washington, DC.

Benthic Macroinvertebrates

Benthic macroinvertebrates were collected to provide a qualitative description of the community composition at each sampling site (Kazyak 1996). Sampling was conducted during the spring index period. Benthic community data were collected for the purpose of calculating biological metrics, such as those described in EPA's Rapid Bioassessment Protocols (Plafkin et al. 1989), and use as an indicator of biological integrity for Maryland streams.

At each segment, a 600 micron mesh "D" net was used to collect organisms from habitats likely to support the greatest taxonomic diversity. A riffle area was preferred, but other habitats were also sampled using a variety of techniques including kicking, jabbing, and gently rubbing hard surfaces by hand to dislodge organisms. If available, other habitat types were sampled, including rootwads, woody debris, leaf packs, macrophytes, and undercut banks. Each jab covered one square foot, and a total of approximately 2.0 m² (20 square feet) of combined substrates was sampled and preserved in 70% ethanol. In the laboratory, the preserved sample was transferred to a gridded pan and organisms were picked from randomly selected grid cells until the cell that contained the 100th individual (if possible) was completely picked. Some samples had fewer than 100 individuals. The benthic macroinvertebrates were identified to genus, or lowest practicable taxon, in the laboratory.

Index of Biotic Integrity

Sites were evaluated using both the fish (F-IBI) and benthic macroinvertebrate (B-IBI) IBIs developed for the MBSS (for detailed methods, see Roth et al. 1997 and Stribling et al. 1998). IBI scores for the MBSS are

determined by comparing the fish or benthic macroinvertebrate assemblages at each site to those found at minimally impacted reference sites. Three separate formulations were employed for the fish IBI, one for each of three distinct geographic areas: Coastal Plain, Eastern Piedmont, and Highland. The two formulations used for the benthic IBI cover the Coastal Plain and non-Coastal Plain regions. Individual metrics for the IBI are scored 1, 3, or 5, based on comparison with the distribution of metric values at reference sites. For either the individual metrics or total IBI, a score of 3 or greater is considered comparable to reference site conditions, while scores falling below this threshold differ significantly from the reference conditions. Scores for the MBSS IBIs are calculated as the mean of the individual metric scores and therefore range from 1 to 5. Some other programs have used a similar approach (e.g., Weisberg et al. 1997), while others have instead computed the IBI as the total of individual metric scores. For example, Karr et al. (1986) calculated IBI as the sum of 12 metric scores, with totals ranging from 12 to 60 points.

Reptiles and Amphibians

At each sample segment, reptiles and amphibians were identified and the presence of observed species was recorded during the summer index period. A search of the riparian area was conducted within 5 meters of the stream on both sides of the 75-meter segment. Any reptiles and amphibians collected during the electrofishing of the stream segment were also included in the species list. Individuals were identified to species when possible. Voucher specimens and individuals not positively identifiable in the field were retained for examination in the laboratory and confirmation by herpetologists at the Smithsonian Institution, Washington, DC, or Towson University, Towson, Maryland.

Physical Habitat

Habitat assessments were conducted at all stream segments as a means of assessing the importance of physical habitat to the biological integrity and fishability of freshwater streams in Maryland. Procedures for habitat assessments (Kazyak 1996) were derived from two currently used methodologies: EPA's Rapid

Bioassessment Protocols (RBPs) (Plafkin et al. 1989), as modified by Barbour and Stribling (1991), and the Ohio EPA's Qualitative Habitat Evaluation Index (QHEI) (Ohio EPA 1987, Rankin 1989). A number of characteristics (instream habitat, epifaunal substrate, velocity/depth diversity, pool/glide/eddy quality, riffle/run quality, channel alteration, bank stability, embeddedness, channel flow status, and shading) were assessed qualitatively, based on visual observations within each 75-meter sample segment. Riparian zone vegetation width was estimated to the nearest meter, up to 50 meters from the stream. Additional observations of the surrounding area were used to assign ratings for aesthetic value (based on visible signs of human refuse at a site) and remoteness (based on distance from the nearest road, accessibility, and evidence of human activity). Also recorded were the presence or absence of various stream features including substrate types, various morphological characteristics, beaver ponds, point sources, and stream channelization. Local land uses visible from the stream segment and riparian vegetation type were also noted. Several additional physical characteristics were measured quantitatively to further characterize the habitat for each segment (see Kazyak 1996 for details). Quantitative measurements of the segment included maximum depth, stream gradient, velocity, thalweg depth, number of functional rootwads, number of functional large woody debris, wetted width, sinuosity, and overbank flood height. A velocity/depth profile was measured or other data were collected to enable calculation of discharge.

Physical Habitat Index

The Physical Habitat Index (PHI) was developed using MBSS data from 1994 to 1997 (Hall et al. 1999). As was the case in development of the fish and benthic IBIs, the conceptual approach was based on evaluating the relative importance (discriminatory power) of individual metrics and combinations of metrics explaining natural differences in streams throughout Maryland. These metrics were derived from both quantitative and qualitative habitat data collected during the summer index period. Based on analyses conducted for both fish IBI (Roth et al. 1998) and benthic macroinvertebrate IBI (Stribling et al. 1998) development in Maryland, the State was divided into two regions: the Coastal Plain and non-Coastal Plain.

The resulting index was then adjusted to a centile scale that rated each sample segment as follows: Good - 72 to 100; Fair - 42 to 71.9; Poor - 12 to 41.9; and Very Poor - 0 to 11.9.

Water Chemistry

During the spring index period, water samples were collected at each site for analysis of pH, acid neutralizing capacity (ANC), conductivity, sulfate, nitrate-nitrogen, and dissolved organic carbon (DOC). These variables describe basic water quality conditions with an emphasis on factors related to acidic deposition.

Grab samples were collected in one-liter bottles for analysis of all analytes except pH. Water samples for pH were collected with 60 ml syringes, which allowed purging of air bubbles to minimize changes in carbon dioxide content (EPA 1987). Samples were stored on wet ice and shipped on wet ice to the analytical laboratory within 48 hours. Laboratory analyses were carried out by the University of Maryland's Appalachian Laboratory in Frostburg.

Chemical analysis of water samples followed standard methods described in EPA's Handbook of Methods for Acid Deposition Studies (EPA 1987). EPA protocols were followed, except that ANC sample volume was reduced to 40 ml to ease handling. Routine daily quality control (QC) checks included processing duplicate, blank, and calibration samples according to EPA guidelines for each analyte. Field duplicates were taken at 5% of all sites. Routine QC checks helped to identify and correct errors in sampling routines or instrumentation at the earliest possible stage.

During the summer index period, in situ measurements of dissolved oxygen (DO), pH, temperature, and conductivity were collected at each site to further characterize existing water quality conditions that might influence biological communities. Measurements were made at an undisturbed section of the segment, usually in the middle of the stream channel, using electrode probes. Instruments were calibrated daily and calibration logbooks were maintained to document instrument performance.

Recognizing that water temperature is an important factor affecting stream condition, but one that varies

daily and seasonally, temperature loggers were deployed at 220 sites in five basins during 1997. The basins sampled were: the Choptank, Susquehanna, Potomac Washington Metro, Patuxent, and Pocomoke. Onset Computer Corporation Optic Stowaway temperature loggers were anchored in each site during the summer index period. Water temperature was recorded every 15 minutes from June 15 until mid-September.

Mussels

During the summer index period, freshwater mussels were sampled qualitatively by examining each 75-meter stream segment for their presence. Mussels were identified to species, their presence recorded, and subsequently released. Species not positively identifiable in the field were retained for confirmation by U.S. Geological Survey (USGS) Biological Resources Division staff.

Aquatic Vegetation

Aquatic vegetation was sampled qualitatively by examining each 75-meter segment for the presence of aquatic plants. Plants were identified to species and their presence recorded for each site. While the primary objective was to document the presence of submerged aquatic vegetation (SAV), emergent and floating aquatic vegetation was also recorded when encountered. Species not positively identifiable in the field were retained for laboratory examination and confirmation by MDNR's staff expert on SAV. Due to the difficulty in long-term preservation, no permanent vouchers of aquatic vegetation were retained.

Data Management

All crews used standardized pre-printed data forms developed for the Survey to ensure that all data for each sampling segment were recorded and standard units of measure were used (Kazyak 1996). Using standard data forms facilitated data entry and minimized transcription error. The field crew leader and a second reviewer checked all data sheets for completeness and legibility before leaving each sampling location. Original data sheets were sent to the Data Management Officer for further review and data entry, while copies were retained by the field crews.

A custom database application, in which the input module was designed to match each of the field data sheets, was used for data entry. Data were independently entered into two databases and compared using a computer program as a quality-control procedure. Differences between the two databases were resolved from original data sheets or through discussions with field crew leaders.

**THIS PAGE INTENTIONALLY
LEFT BLANK**

Maryland Biological Stream Survey Data

COUNTY SUMMARY

A total of 73 quantitative sites were sampled in Harford county by MBSS sampling crews during 1994-1997 (Table 1; Figure 2). In addition, the fish assemblage was characterized on a presence/absence basis at 55 sites to provide a more complete picture of fish species distributions. Appendix A provides a summary of the types of data available for each of the sites sampled.

Species Highlights

A total of 48 fish species were collected in the small to mid-sized streams (Table 2); this number ties for a ranking of fourth in the state. Like most other areas in the state, nearly all (98%) of the sites sampled contained fish. In contrast, the 172 genera of benthic macroinvertebrates found in Harford county ties this area for a ranking of ninth best among Maryland counties for benthic diversity (Table 3).

Blacknose dace, creek chub, and rosyside dace were the most commonly encountered fish species (present at 90% of sites sampled) in Harford county (Table 2). Brook trout, a temperature-sensitive species, was present at only 7% of the sites sampled. It is likely that historical agricultural land use and increasing urbanization (and subsequent increase in runoff from impervious surfaces) are the reasons that brook trout are now rare in the county and undoubtedly declining. Logperch, a rare fish species in Maryland, was collected at a single site.

Sixteen species of reptiles and amphibians were found in or near Harford county streams (Table 4), tying the county for a ranking of thirteenth in the state. No state or federally listed species were collected.

Ecological Health

The overall ecological health of Harford county's headwater streams can best be described as Fair. The average F-IBI score was 3.75 (high end of the Fair range) and the average B-IBI score among sites was 3.0 (Fair category, but nearly in the Poor). Based on F-IBI and B-IBI scores from individual sites, one of the highest rated streams in the county is Deep Creek. The worst streams include the Middle Branch of Romney Creek and Herring Run (Table 6).

Physical Habitat

Physical habitat in Harford County was rated as Fair by the Physical Habitat Index. Values ranged from 5.2 to 96.9, with an average score of 66.5 (upper end of the Fair range, ranking second among counties in the state) (Table 6; Figure 5). Other noteworthy points about Harford County streams include instream habitat and epifaunal substrate scores, with an average rating of 14 and 13, respectively, ranking second in the state.

Nitrate-Nitrogen

Nitrate-nitrogen values at sites sampled in Harford county averaged 3.02 mg/L, making Harford the eighth worst county in the state. The stream with the lowest nitrate values in the county was the Middle Branch of Romney Creek while the streams with the highest nitrate values included Jack's Hole and Big Branch (Table 7). In no stream was the EPA limit for drinking water (10 mg/L) exceeded.

Table 1. Site information and land use data collected at Maryland Biological Stream Survey sites in Harford County, 1994-1997. Basin abbreviations are as follows: BU - Bush River; GU - Gunpowder River; SQ - Lower Susquehanna River.

Site	Latitude	Longitude	Stream Name	Basin	Order	Catchment Acres	% Urban	% Agric.	% Forest
BA-P-042-116-96	39.5920	76.5560	Little Gunpowder Fls	GU	1	1570.21	0.02	68.17	31.76
BA-P-215-305-96	39.5360	76.5200	Little Gunpowder R	GU	3	12691.01	0.33	70.14	28.56
HA-N-003-204-96	39.4560	76.1680	Romney Cr Middle Br	BU	2	2482.40	48.50	14.55	20.50
HA-N-009-105-96	39.4420	76.3440	Un Str	BU	1	101.73	26.42	23.41	50.17
HA-N-018-103-96	39.5670	76.2020	Swan Cr	BU	1	184.07	2.77	80.04	15.71
HA-N-036-206-96	39.5320	76.1670	Carsins Run	BU	2	6738.48	5.12	48.88	44.73
HA-N-040-307-96	39.4570	76.3240	Winters Run	BU	3	33186.82	6.83	53.01	39.88
HA-N-052-202-96	39.5230	76.1260	Gasheys Run	BU	2	2626.66	5.61	40.85	53.03
HA-N-067-111-96	39.4990	76.1980	Cranberry Run	BU	1	1415.06	8.68	34.65	56.47
HA-N-068-301-96	39.5070	76.3100	Bynum Run	BU	3	8327.75	19.56	53.07	27.21
HA-N-068-308-96	39.5040	76.3080	Bynum Run	BU	3	8422.34	19.34	52.88	27.62
HA-N-086-201-96	39.5290	76.2650	Broad Run	BU	2	1972.04	5.00	68.47	26.16
HA-N-092-304-96	39.4690	76.3370	Winters Run	BU	3	30987.84	7.00	53.60	39.10
HA-N-099-305-96	39.4790	76.2810	Bynum Run	BU	3	13323.18	15.69	52.05	31.50
HA-P-001-205-96	39.5780	76.4390	East Br	BU	2	4182.93	1.63	55.67	42.12
HA-P-005-206-97	39.7070	76.4830	Big Br	SQ	2	2575.65	0.19	77.86	21.83
HA-P-008-1-94	39.7016	76.2836	Deep Cr	SQ	2	3456.90	0.05	57.92	41.81
HA-P-008-3-94	39.6919	76.2606	Deep Cr	SQ	2	4445.50	0.17	52.80	46.78
HA-P-010-103-97	39.5970	76.4080	South Stirrup Run	SQ	1	976.44	0.56	62.54	36.91
HA-P-012-205-97	39.6440	76.3580	Mine Br	SQ	2	1986.32	0.33	63.03	36.35
HA-P-013-101-97	39.5940	76.1360	Herring Run	SQ	1	868.27	4.08	42.51	53.41
HA-P-024-116-97	39.6610	76.3540	Un Trib To Broad Cr	SQ	1	56.72	0.00	80.84	19.16
HA-P-033-203-97	39.6630	76.4710	Little Deer Cr	SQ	2	8115.48	1.04	66.38	32.02
HA-P-033-212-97	39.6610	76.4470	Little Deer Cr	SQ	2	8918.23	0.95	66.36	32.18
HA-P-035-208-97	39.6800	76.5570	Plumtree Br	SQ	2	23233.82	1.60	67.33	30.39
HA-P-037-104-97	39.6830	76.4080	Jacks Hole	SQ	1	198.71	0.00	87.37	12.63
HA-P-038-106-96	39.5240	76.3690	Heavenly Waters	BU	1	683.89	17.26	44.23	38.51
HA-P-041-1-94	39.6742	76.5259	Deer Cr	SQ	1	422.50	0.07	71.62	28.31
HA-P-041-2-94	39.6683	76.5237	Deer Cr	SQ	1	246.80	0.00	85.90	14.10
HA-P-062-207-96	39.5520	76.3430	Bynum Run	BU	2	2745.07	15.40	56.70	27.74
HA-P-068-114-97	39.7170	76.4920	Big Br	SQ	1	1115.71	0.27	78.89	20.75
HA-P-071-1-94	39.6855	76.3671	Deep Cr	SQ	3	7383.30	0.79	75.09	23.42
HA-P-071-2-94	39.6784	76.3589	Deep Cr	SQ	3	8041.00	0.73	75.27	23.36
HA-P-071-318-97	39.6830	76.3640	Broad Cr	SQ	3	7443.85	0.78	75.21	23.31
HA-P-078-4-94	39.5992	76.3644	Cabbage Run Br	SQ	1	841.40	0.17	87.93	11.79
HA-P-078-6-94	39.5987	76.3688	Cabbage Run Br	SQ	1	609.00	0.23	93.27	6.45
HA-P-087-1-94	39.6142	76.3412	Stout Bottle Br	SQ	2	3108.50	1.92	73.70	24.10
HA-P-087-207-97	39.6080	76.3480	Cabbage Run Br	SQ	2	2782.70	2.13	76.35	21.21
HA-P-087-2-94	39.6091	76.3465	Stout Bottle Br	SQ	2	2800.50	2.14	75.97	21.59
HA-P-099-102-97	39.6470	76.2220	Holland's Br	SQ	1	98.18	0.00	20.71	79.29
HA-P-100-204-97	39.6500	76.4990	Deer Cr	SQ	2	4167.90	0.24	73.17	25.65
HA-P-116-109-96	39.5090	76.4250	Un Trib To Little Gunpowder R	GU	1	548.81	5.28	43.27	51.46
HA-P-128-104-96	39.5530	76.4760	Un Trib To West Br Winters Run	BU	1	563.10	0.73	86.19	13.02
HA-P-131-1-94	39.7076	76.2900	Deep Cr	SQ	2	2786.80	0.05	56.76	43.09
HA-P-131-3-94	39.7063	76.2949	Deep Cr	SQ	2	2703.70	0.05	58.16	41.69
HA-P-133-111-97	39.6310	76.3040	Un Trib To Deer Cr	SQ	1	553.70	0.00	50.31	49.69

Table 1 (cont.). Site information and land use data collected at Maryland Biological Stream Survey sites in Harford County, 1994-1997. Basin abbreviations are as follows: BU - Bush River; GU - Gunpowder River; SQ - Lower Susquehanna River.

Site	Latitude	Longitude	Stream Name	Basin	Order	Catchment Acres	% Urban	% Agric.	% Forest
HA-P-138-302-97	39.6830	76.2800	Broad Cr	SQ	3	18206.78	0.52	62.69	35.60
HA-P-141-302-96	39.5440	76.4240	Winters Run	BU	3	13589.93	1.30	57.85	40.54
HA-P-141-303-96	39.5510	76.4370	Winters Run	BU	3	12622.96	1.24	58.43	40.00
HA-P-142-105-97	39.6530	76.3060	Un Trib To Broad Cr	SQ	1	308.00	0.22	68.30	31.36
HA-P-151-102-96	39.5070	76.3400	Plumtree Run	BU	1	1216.70	46.41	31.57	22.00
HA-P-156-112-97	39.6330	76.4810	Un Trib To Deer Cr	SQ	1	272.53	6.74	52.19	41.07
HA-P-164-306-96	39.5260	76.3170	Bynum Run	BU	3	6902.82	21.82	51.43	26.56
HA-P-174-1-94	39.6623	76.3339	Broad Cr	SQ	3	10445.40	0.56	71.15	27.56
HA-P-174-2-94	39.6610	76.3292	Broad Cr	SQ	3	10489.30	0.56	71.15	27.57
HA-P-174-4-94	39.6610	76.3301	Broad Cr	SQ	3	10482.20	0.56	71.14	27.59
HA-P-176-309-97	39.6700	76.3140	Broad Cr	SQ	3	13500.67	0.44	68.70	29.46
HA-P-178-202-97	39.6960	76.4650	Big Br	SQ	2	4084.09	0.13	72.81	26.92
HA-P-178-209-97	39.6830	76.4600	Big Br	SQ	2	4789.73	0.11	66.39	33.38
HA-P-180-107-97	39.7150	76.4430	Falling Br	SQ	1	1396.05	2.82	86.20	10.73
HA-P-186-308-97	39.6680	76.2950	Broad Cr	SQ	3	15201.21	0.39	66.16	32.18
HA-P-205-1-94	39.6665	76.2802	Broad Cr	SQ	1	487.80	2.58	37.51	59.91
HA-P-205-2-94	39.6690	76.2791	Broad Cr	SQ	1	804.60	2.85	30.44	66.71
HA-P-207-1-94	39.6189	76.3399	Stout Bottle Br	SQ	2	4557.20	1.31	74.89	23.60
HA-P-207-2-94	39.6201	76.3369	Stout Bottle Br	SQ	2	4629.30	1.29	74.97	23.54
HA-P-214-211-97	39.6290	76.2240	Holland Br	SQ	2	1934.47	0.14	54.09	45.68
HA-P-216-1-94	39.6399	76.2609	Hopkins Br	SQ	1	251.50	1.07	33.32	65.61
HA-P-216-2-94	39.6173	76.2325	Hopkins Br	SQ	1	1460.50	0.23	47.19	52.49
HA-P-225-1-94	39.7033	76.3036	Deep Cr	SQ	1	376.20	0.08	71.03	28.71
HA-P-225-2-94	39.7017	76.3051	Deep Cr	SQ	1	356.50	0.08	73.61	26.11
HA-P-244-1-94	39.5904	76.3565	Stout Bottle Br	SQ	1	532.10	7.84	68.59	22.98
HA-P-244-2-94	39.5996	76.3552	Stout Bottle Br	SQ	1	1069.70	5.32	72.70	21.69
HA-P-246-304-96	39.5240	76.5040	Little Gunpowder R	GU	3	15305.43	0.43	65.27	33.38




Figure 1. Land use in Harford County (MOP 1994).




Figure 2. Location of Maryland Biological Stream Survey sites in Harford County, 1994-1997.

Table 2. Percent occurrence of fish species collected at Maryland Biological Stream Survey sites in Harford County, 1994-1997.

Family	Common Name	Scientific Name	Number of Occurrences	Percent Occurrence
Petromyzontidae	least brook lamprey	<i>Lampetra aepyptera</i>	3	4.35
	sea lamprey	<i>Petromyzon marinus</i>	14	20.29
Anguillidae	American eel	<i>Anguilla rostrata</i>	43	62.32
Clupeidae	gizzard shad	<i>Dorosoma cepedianum</i>	1	1.45
Cyprinidae	central stoneroller	<i>Camptostoma anomalum</i>	10	14.49
	goldfish	<i>Carassius auratus</i>	1	1.45
	rosyside dace	<i>Clinostomus funduloides</i>	60	86.96
	satinfin shiner	<i>Cyprinella analostana</i>	17	24.64
	spotfin shiner	<i>Cyprinella spiloptera</i>	1	1.45
	common carp	<i>Cyprinus carpio</i>	1	1.45
	cutlips minnow	<i>Exoglossum maxillingua</i>	47	68.12
	eastern silvery minnow	<i>Hybognathus regius</i>	1	1.45
	common shiner	<i>Luxilus cornutus</i>	37	53.62
	river chub	<i>Nocomis micropogon</i>	30	43.48
	golden shiner	<i>Notemigonus crysoleucas</i>	2	2.90
	comely shiner	<i>Notropis amoenus</i>	2	2.90
	spottail shiner	<i>Notropis hudsonius</i>	8	11.59
	swallowtail shiner	<i>Notropis procne</i>	21	30.43
	rosyface shiner	<i>Notropis rubellus</i>	7	10.14
	bluntnose minnow	<i>Pimephales notatus</i>	15	21.74
	blacknose dace	<i>Rhinichthys atratulus</i>	63	91.30
	longnose dace	<i>Rhinichthys cataractae</i>	42	60.87
	creek chub	<i>Semotilus atromaculatus</i>	64	92.75
	fallfish	<i>Semotilus corporalis</i>	30	43.48
Catostomidae	white sucker	<i>Catostomus commersoni</i>	57	82.61
	creek chubsucker	<i>Erimyzon oblongus</i>	4	5.80
	northern hogsucker	<i>Hypentelium nigricans</i>	36	52.17
Ictaluridae	yellow bullhead	<i>Ameiurus natalis</i>	2	2.90
	brown bullhead	<i>Ameiurus nebulosus</i>	2	2.90
	marginated madtom	<i>Noturus insignis</i>	38	55.07
Esocidae	redfin pickerel ¹	<i>Esox americanus vermiculatus</i>		
Umbridae	eastern mudminnow	<i>Umbra pygmaea</i>	1	1.45
Salmonidae	rainbow trout	<i>Oncorhynchus mykiss</i>	1	1.45
	brown trout	<i>Salmo trutta</i>	9	13.04
	brook trout	<i>Salvelinus fontinalis</i>	5	7.25
Cottidae	mottled sculpin	<i>Cottus bairdi</i>	26	37.68
Centrarchidae	rock bass	<i>Ambloplites rupestris</i>	5	7.25
	banded sunfish ¹	<i>Enneacanthus obesus</i>		
	redbreast sunfish	<i>Lepomis auritus</i>	28	40.58
	green sunfish	<i>Lepomis cyanellus</i>	13	18.84
	pumpkinseed	<i>Lepomis gibbosus</i>	20	28.99
	bluegill	<i>Lepomis macrochirus</i>	24	34.78
	smallmouth bass	<i>Micropterus dolomieu</i>	18	26.09
	largemouth bass	<i>Micropterus salmoides</i>	13	18.84
Percidae	tessellated darter	<i>Etheostoma olmstedi</i>	55	79.71
	banded darter	<i>Etheostoma zonale</i>	1	1.45
	logperch	<i>Percina caprodes</i>	1	1.45
	shield darter	<i>Percina peltata</i>	7	10.14
None			2	2.90

¹ Qualitative Sites




Figure 3. Stream ecological conditions based on the Fish Index of Biotic Integrity (F-IBI) at Maryland Biological Stream Survey sites in Harford County, 1994-1997.

Table 3. Tolerance Value (TV)¹, Functional Feeding Group (FFG), Habit, and Percent Occurrence of benthic macroinvertebrate taxa² collected at Maryland Biological Stream Survey sites in Harford County, 1994-1997. Abbreviations of habits are as follows: bu - burrower, cn - clinger, cb - climber, sp - sprawler, dv - diver, and sk - skater.

Class	Order	Family	Genus	TV	FFG	Habit	Percent Occurrence
Enopla	Hoplonemertea	Tetrastemmatidae	<i>Prostoma</i> Sp.		Predator		1.67
Turbellaria	Tricladida	Planariidae	<i>Cura</i> Sp.			sp	1.67
Oligochaeta	Lumbriculida	Lumbriculidae		10	Collector	bu	13.33
Oligochaeta	Tubificida	Enchytraeidae		10	Collector	bu	1.67
		Naididae		10	Collector	bu	5.00
		Tubificidae		10	Collector	cn	3.33
			<i>Limnodrilus</i> Sp.	10	Collector	cn	3.33
Gastropoda	Basommatophora	Ancylidae	<i>Fissia</i> Sp.	7	Scraper	cb	1.67
		Lymnaeidae	<i>Pseudosuccinea</i> Sp.	6	Collector	cb	1.67
		Physidae	<i>Physella</i> Sp.	8	Scraper	cb	6.67
		Planorbidae	<i>Planorbella</i> Sp.	7	Scraper	cb	1.67
Gastropoda	Mesogastropoda	Pleuroceridae	<i>Leptaxis</i> Sp.		Scraper	cb	1.67
Pelecypoda	Veneroida	Sphaeriidae	<i>Psidium</i> Sp.	8	Filterer	bu	1.67
			<i>Sphaerium</i> Sp.	8	Filterer	bu	1.67
Malacostraca	Amphipoda	Crangonyctidae	<i>Crangonyx</i> Sp.	4	Collector	sp	6.67
		Gammaridae	<i>Gammarus</i> Sp.	6	Shredder	sp	10.00
			<i>Stygonectes</i> Sp.	6	Shredder	sp	3.33
						sp	1.67
Malacostraca	Decapoda	Cambaridae		6	Shredder	sp	1.67
			<i>Cambarus</i> Sp.	6	Collector	sp	3.33
		Palaemonidae	<i>Palaemonetes</i> Sp.	7		sp	1.67
Malacostraca	Isopoda	Asellidae	<i>Caecidotea</i> Sp.	8	Collector	sp	1.67
Insecta	Collembola						1.67
Insecta	Ephemeroptera	Ameletidae	<i>Ameletus</i> Sp.	0	Collector	sw, cb	18.33
		Baetidae			Collector	sw, cn	8.33
			<i>Acentrella</i> Sp.	4	Collector	sw, cn	3.33
			<i>Acerpenna</i> Sp.	4	Collector	sw, cn	10.00
			<i>Baetis</i> Sp.	6	Collector	sw, cb, cn	20.00
			<i>Centroptilum</i> Sp.	2	Collector	sw, cn	3.33
		Caenidae	<i>Caenis</i> Sp.	7	Collector	sp	6.67
		Ephemerellidae				cn, sp, sw	1.67
			<i>Drunella</i> Sp.	1	Scraper	cn, sp	1.67
			<i>Ephemerella</i> Sp.	2	Collector	cn, sw	70.00
			<i>Eurylophella</i> Sp.	4	Scraper	cn, sp	45.00
			<i>Satella</i> Sp.	2	Collector	cn	11.67
		Heptageniidae	<i>Epeorus</i> Sp.	0	Scraper	cn	40.00
			<i>Heptagenia</i> Sp.	4	Scraper	cn, sw	5.00
			<i>Stenacron</i> Sp.	4	Collector	cn	6.67
			<i>Stenonema</i> Sp.	4	Scraper	cn	56.67
		Isonychiidae	<i>Isonychia</i> Sp.	2	Filterer	sw, cn	16.67
		Leptophlebiidae			Collector	sw, cn	1.67
			<i>Habrophlebia</i> Sp.		Collector	sw, cn, sp	1.67
			<i>Leptophlebia</i> Sp.	4	Collector	sw, cn, sp	3.33
			<i>Paraleptophlebia</i> Sp.	2	Collector	sw, cn, sp	21.67
Insecta	Odonata	Aeshnidae	<i>Boyeria</i> Sp.	2	Predator	cb, sp	5.00
		Calopterygidae	<i>Calopteryx</i> Sp.	6	Predator	cb	1.67
		Coenagrionidae	<i>Argia</i> Sp.	8	Predator	cn, cb, sp	8.33

Table 3 (cont.). Tolerance Value (TV)¹, Functional Feeding Group (FFG), Habit, and Percent Occurrence of benthic macroinvertebrate taxa² collected at Maryland Biological Stream Survey sites in Harford County, 1994-1997. Abbreviations of habits are as follows: bu - burrower, cn - clinger, cb - climber, sp - sprawler, dv - diver, and sk - skater.

Class	Order	Family	Genus	TV	FFG	Habit	Percent Occurrence
Insecta	Plecoptera	Gomphidae			Predator	bu	3.33
			<i>Arigomphus</i> Sp.	42	Predator	bu	1.67
			<i>Gomphus</i> Sp.	5	Predator	bu	1.67
			<i>Stylogomphus</i> Sp.		Predator	bu	5.00
		Capniidae			Shredder	sp, cn	1.67
			<i>Allocapnia</i> Sp.	3	Shredder	cn	1.67
			<i>Paracapnia</i> Sp.	1	Shredder	-	1.67
		Chloroperlidae			Predator	cn	1.67
			<i>Sweltsa</i> Sp.		Predator	cn	3.33
		Leuctridae			Shredder	sp, cn	6.67
			<i>Leuctra</i> Sp.	0	Shredder	cn	10.00
		Nemouridae			Shredder	sp, cn	3.33
			<i>Amphinemura</i> Sp.	3	Shredder	sp, cn	31.67
			<i>Ostrocerca</i> Sp.		Shredder	sp, cn	1.67
			<i>Prostoia</i> Sp.		Shredder	sp, cn	28.33
		Peltoperlidae			Shredder	cn, sp	1.67
		Perlidae			Predator	cn	6.67
			<i>Acroneuria</i> Sp.	0	Predator	cn	15.00
			<i>Eccoptura</i> Sp.		Predator	cn	6.67
		Perlodidae			Predator	cn	10.00
			<i>Clasperla</i> Sp.	1	Predator	cn	1.67
			<i>Diploperla</i> Sp.		Predator	cn	3.33
			<i>Isoperla</i> Sp.	2	Predator	cn, sp	3.33
		Pteronarcyidae	<i>Pteronarcys</i> Sp.	2	Shredder	cn, sp	1.67
Insecta	Megaloptera	Taeniopterygidae	<i>Strophopteryx</i> Sp.		Shredder	sp, cn	3.33
		Corydalidae	<i>Corydalus</i> Sp.	5	Predator	cn, cb	11.67
			<i>Nigronia</i> Sp.	0	Predator	cn, cb	16.67
Insecta	Trichoptera	Sialidae	<i>Sialis</i> Sp.	4	Predator	bu, cb, cn	5.00
		Glossosomatidae	<i>Glossosoma</i> Sp.	0	Scraper	cn	1.67
					Filterer	cn	1.67
		Hydropsychidae	<i>Cheumatopsyche</i> Sp.	5	Filterer	cn	58.33
			<i>Diplectrona</i> Sp.	2	Filterer	cn	30.00
			<i>Hydropsyche</i> Sp.	6	Filterer	cn	55.00
		Limnephilidae			Shredder	cb, sp, cn	1.67
			<i>Goera</i> Sp.		Scraper	cn	1.67
			<i>Ironoquia</i> Sp.	3	Shredder	sp	1.67
			<i>Pycnopsyche</i> Sp.	4	Shredder	sp, cb, cn	6.67
		Odontoceridae	<i>Psilotreta</i> Sp.	0	Scraper	sp	1.67
		Philopotamidae	<i>Chimarra</i> Sp.	4	Filterer	cn	18.33
			<i>Dolophilodes</i> Sp.	0	Filterer	cn	8.33
		Polycentropodidae				cn	1.67
			<i>Polycentropus</i> Sp.	5	Filterer	cn	8.33
		Psychomyiidae	<i>Psychomyia</i> Sp.	2	Collector	cn	5.00
		Rhyacophilidae	<i>Rhyacophila</i> Sp.	1	Predator	cn	23.33
		Uenoidae	<i>Neophylax</i> Sp.	3	Scraper	cn	20.00
Insecta	Coleoptera	Dryopidae	<i>Helichus</i> Sp.	5	Scraper	cn	8.33

Table 3 (cont.). Tolerance Value (TV)¹, Functional Feeding Group (FFG), Habit, and Percent Occurrence of benthic macroinvertebrate taxa² collected at Maryland Biological Stream Survey sites in Harford County, 1994-1997. Abbreviations of habits are as follows: bu - burrower, cn - clinger, cb - climber, sp - sprawler, dv - diver, and sk - skater.

Class	Order	Family	Genus	TV	FFG	Habit	Percent Occurrence
Insecta	Diptera	Dytiscidae	<i>Agabus</i> Sp.	5	Predator	sw, dv	3.33
			<i>Hydroporus</i> Sp.	5	Predator	sw, cb	1.67
		Elmidae		5	Collector	cn	1.67
			<i>Dubiraphia</i> Sp.	6	Scraper	cn, cb	8.33
			<i>Optioservus</i> Sp.	4	Scraper	cn	40.00
			<i>Oulimnius</i> Sp.	2	Scraper	cn	35.00
			<i>Promoresia</i> Sp.	2	Scraper	cn	1.67
			<i>Stenelmis</i> Sp.	6	Scraper	cn	15.00
		Gyrinidae	<i>Dineutus</i> Sp.	4	Predator	sw, dv	1.67
		Haliplidae	<i>Peltodytes</i> Sp.	5	Shredder	cb, cn	1.67
		Hydrophilidae	<i>Hydrobius</i> Sp.	5	Collector	cb, cn, sp	1.67
		Psephenidae	<i>Ectopria</i> Sp.	5	Scraper	cn	3.33
			<i>Psephenus</i> Sp.	4	Scraper	cn	10.00
		Ptilodactylidae	<i>Anchytarsus</i> Sp.	4	Shredder	cn	5.00
		Athericidae	<i>Atherix</i> Sp.	2	Predator	sp, bu	1.67
		Ceratopogonidae			Predator	sp, bu	1.67
			<i>Bezzia</i> Sp.	6	Predator	bu	5.00
			<i>Ceratopogon</i> Sp.	6	Predator	sp, bu	1.67
			<i>Probezzia</i> Sp.	6	Predator	bu	6.67
		Chironomidae	<i>Ablabesmyia</i> Sp.	8	Predator	sp	3.33
			<i>Apsectrotanytus</i> Sp.	5	Predator	bu, sp	1.67
			<i>Brillia</i> Sp.	5	Shredder	bu, sp	6.67
			<i>Cladotanytarsus</i> Sp.	7	Filterer	-	1.67
			<i>Conchapelopia</i> Sp.	6	Predator	sp	31.67
			<i>Corynoneura</i> Sp.	7	Collector	sp	5.00
			<i>Cricotopus</i> Sp.	7	Shredder	cn, bu	6.67
			<i>Cricotopus</i> /				
			<i>Orthocladius</i> Sp.		Shredder		63.33
			<i>Cryptochironomus</i> Sp.	8	Predator	sp, bu	3.33
			<i>Diamesa</i> Sp.	5	Collector	sp	25.00
			<i>Dicrotendipes</i> Sp.	10	Collector	bu	1.67
			<i>Diplocladius</i> Sp.	7	Collector	sp	1.67
			<i>Endobironomus</i> Sp.	10	Shredder	cn	1.67
			<i>Eukiefferiella</i> Sp.	8	Collector	sp	35.00
			<i>Heleniella</i> Sp.		Predator	sp	1.67
			<i>Heterotrissocladius</i> Sp.		Collector	sp, bu	3.33
			<i>Hydrobaenus</i> Sp.	8	Scraper	sp	1.67
			<i>Micropectra</i> Sp.	7	Collector	cb, sp	10.00
			<i>Microtendipes</i> Sp.	6	Filterer	cn	10.00
			<i>Nanocladius</i> Sp.	3	Collector	sp	5.00
			<i>Orthocladinae</i> A Sp.		Collector		21.67
			<i>Orthocladius</i> Sp.	6	Collector	sp, bu	35.00
			<i>Pagastia</i> Sp.	1	Collector	-	11.67
			<i>Parametriocnemus</i> Sp.	5	Collector	sp	36.67
			<i>Paratanytarsus</i> Sp.	6	Collector	sp	1.67
			<i>Pentaneura</i> Sp.	6	Predator	sp	1.67
			<i>Polypedilum</i> Sp.	6	Shredder	cb, cn	35.00

Table 3 (cont.). Tolerance Value (TV)¹, Functional Feeding Group (FFG), Habit, and Percent Occurrence of benthic macroinvertebrate taxa² collected at Maryland Biological Stream Survey sites in Harford County, 1994-1997. Abbreviations of habits are as follows: bu - burrower, cn - clinger, cb - climber, sp - sprawler, dv - diver, and sk - skater.

Class	Order	Family	Genus	TV	FFG	Habit	Percent Occurrence
			<i>Rheocricotopus</i> Sp.	6	Collector	sp	8.33
			<i>Rheotanytarsus</i> Sp.	6	Filterer	cn	20.00
			<i>Sublettea</i> Sp.		Collector	-	5.00
			<i>Sympotthastia</i> Sp.	2	Collector	sp	10.00
			<i>Tanytarsus</i> Sp.	6	Filterer	cb, cn	15.00
			<i>Thienemanniella</i> Sp.	6	Collector	sp	11.67
			<i>Thienemannimyia</i> Sp.		Predator	sp	10.00
			<i>Trissopelopia</i> Sp.		Predator	sp	10.00
			<i>Tvetenia</i> Sp.	5	Collector	sp	6.67
			DIAMESINAE		Collector		1.67
			TANYTARSINI		Collector		3.33
			<i>Zavrelimyia</i> Sp.	8	Predator	sp	5.00
		Dolichopodidae		4	Predator	sp, bu	1.67
		Empididae			Predator	sp, bu	1.67
			<i>Chelifera</i> Sp.		Predator	sp, bu	11.67
			<i>Clinocera</i> Sp.		Predator	cn	16.67
			<i>Hemerodromia</i> Sp.	6	Predator	sp, bu	36.67
		Muscidae	<i>Limnophora</i> Sp.		Predator	bu	1.67
		Simuliidae		7	Filterer	cn	1.67
			<i>Prosimulium</i> Sp.	7	Filterer	cn	53.33
			<i>Simulium</i> Sp.	7	Filterer	cn	16.67
			<i>Stegopterna</i> Sp.	7	Filterer	cn	11.67
		Tabanidae	<i>Chrysops</i> Sp.	7	Predator	sp, bu	5.00
		Tipulidae			Predator	bu, sp	1.67
			<i>Antocha</i> Sp.	5	Collector	cn	40.00
			<i>Dicranota</i> Sp.	4	Predator	sp, bu	3.33
			<i>Hexatoma</i> Sp.	4	Predator	bu, sp	1.67
			<i>Limonia</i> Sp.	6	Shredder	bu, sp	3.33
			<i>Ormosia</i> Sp.		Collector	bu	1.67
			<i>Pseudolimnophila</i> Sp.	2	Predator	bu	3.33
			<i>Rhabdomastix</i> Sp.			bu	1.67
			<i>Tipula</i> Sp.	4	Shredder	bu	15.00

¹ Tolerance values are on a 0 (extremely sensitive) to 10 (tolerant) scale.

² Taxa not identified to genus are presented in capital letters. Subfamily - Diamesinae; Tribe - Tanytarsini.




Figure 4. Stream ecological conditions based on the Benthic Macroinvertebrate Index of Biotic Integrity (B-IBI) at Maryland Biological Stream Survey sites in Harford County, 1994-1997.

Table 4. Percent occurrence of reptile and amphibian species collected at Maryland Biological Stream Survey sites in Harford County, 1994-1997.

Family	Common Name	Scientific Name	Number of Occurrences	Percent Occurrence
Plethodontidae	northern dusky salamander	<i>Desmognathus f. fuscus</i>	15	21.74
	northern two-lined salamander	<i>Eurycea bislineata</i>	31	44.93
	red salamander	<i>Pseudotriton ruber</i>	4	5.80
	redback salamander	<i>Plethodon cinereus</i>	4	5.80
Bufonidae	American toad	<i>Bufo americanus</i>	9	13.04
	Fowler's toad	<i>Bufo woodhousii fowleri</i>	3	4.35
Ranidae	bullfrog	<i>Rana catesbeiana</i>	24	34.78
	green frog	<i>Rana clamitans melanota</i>	17	24.64
	pickerel frog	<i>Rana palustris</i>	21	30.43
	wood frog	<i>Rana sylvatica</i>	2	2.90
Chelydridae	common snapping turtle	<i>Chelydra serpentina</i>	11	15.94
Emydidae	eastern box turtle	<i>Terrapene c. carolina</i>	5	7.25
	wood turtle	<i>Clemmys insculpta</i>	1	1.45
Colubridae	black rat snake	<i>Elaphe o. obsoleta</i>	1	1.45
	eastern garter snake	<i>Thamnophis s. sirtalis</i>	2	2.90
	northern water snake	<i>Nerodia s. sipedon</i>	6	8.70
None			5	7.25

Table 5. Physical habitat data for Maryland Biological Stream Survey sites in Harford County, 1994-1997.

	Instream Habitat ¹	Velocity/Depth Diversity ¹			Riffle Quality ¹	Percent Shading ¹		Number of Woody Debris		Percent Channel Flow ¹		Bank Stability ¹		Aesthetic Rating ¹	
Site		Epifaunal Substrate ¹		Pool Quality ¹		Percent Embeddedness ¹		Maximum Depth (cm) ¹		Number of Rootwads		Channel Alteration ¹		Riparian Width (m) ¹	
BA-P-042-116-96	12	11	13	15	15	70	75	80	4	6	75	5	9	0	16
BA-P-215-305-96	15	14	16	11	16	30	65	84	4	0	85	10	7	50	17
HA-N-009-105-96	8	9	12	14	7	75	70	61	2	6	60	8	11	5	5
HA-N-018-103-96	10	2	5	7	6	90	80	44	0	5	60	7	6	50	16
HA-N-036-206-96	10	15	8	14	3	40	85	73	1	3	80	14	6	15	7
HA-N-040-307-96	10	11	13	13	16	50	40	97	1	2	99	16	12	10	14
HA-N-052-202-96	18	18	17	16	16	30	70	81	5	3	40	9	5	50	15
HA-N-067-111-96	13	10	15	14	13	40	80	120	5	4	50	6	8	50	16
HA-N-068-301-96	18	17	16	16	16	35	70	145	0	6	95	14	14	3	13
HA-N-068-308-96	16	17	10	10	17	25	75	44	0	0	98	16	16	8	13
HA-N-086-201-96	17	16	16	14	16	40	70	56	0	4	60	6	8	50	16
HA-N-092-304-96	17	17	14	12	18	35	40	53	1	0	90	15	17	3	17
HA-N-099-305-96	17	13	18	15	17	40	60	77	0	0	99	10	9	50	13
HA-P-001-205-96	17	14	16	15	15	50	70	79	4	4	90	8	15	50	15
HA-P-005-206-97	17	16	14	17	13	20	30	73	1	1	90	14	8	35	7
HA-P-008-1-94	15	13	11	8	15	30	70	83	6		95	10	8	50	16
HA-P-008-3-94	16	19	9	14	16	10	85	38	0		90	15	18	50	19
HA-P-010-103-97	16	18	11	16	8	10	94	52	1	3	80	9	14	50	16
HA-P-012-205-97	17	16	17	16	16	30	70	86	3	2	85	13	9	15	15
HA-P-013-101-97	17	17	7	13	11	10	95	28	1	1	70	15	17	15	15
HA-P-033-203-97	19	17	19	17	18	15	80	110	1	1	95	16	15	0	15
HA-P-033-212-97	12	12	17	19	5	40	40	96	4	3	99	16	12	3	13
HA-P-035-208-97	18	15	8	13	16	40	95	48	0	1	85	15	16	50	19
HA-P-037-104-97	15	17	8	15	10	30	98	42	4	3	90	10	9	5	13
HA-P-038-106-96	14	12	15	15	14	60	80	97	1	1	85	15	17	30	15
HA-P-041-1-94	17	14	10	9	12	25	95	25	3		70	15	16	50	17
HA-P-041-2-94	5	9	4	3	6	75	95	37	2		50	6	7	7	16
HA-P-062-207-96	8	4	6	16	0	90	60	118	12	3	100	3	5	35	10
HA-P-068-114-97	16	13	13	18	14	15	90	105	0	1	80	9	12	20	9
HA-P-071-1-94	16	15	18	17	17	20	80	85	1		85	18	19	5	15
HA-P-071-2-94	16	15	15	15	17	20	20	130	4		90	16	11	0	14
HA-P-071-318-97	16	17	18	16	16	40	30	95	2	3	85	13	8	23	9
HA-P-078-4-94	16	17	17	16	14	40	20	62	1		95	14	10	0	17
HA-P-078-6-94	16	13	16	16	15	65	50	95	1		80	13	14	50	8
HA-P-087-1-94	12	11	14	13	14	35	40	60	0		60	7	6	0	16
HA-P-087-2-94	17	18	18	19	18	25	60	120	4		95	15	14	10	17

Table 5 (cont.). Physical habitat data for Maryland Biological Stream Survey sites in Harford County, 1994-1997.

	Instream Habitat ¹	Velocity/Depth Diversity ¹			Riffle Quality ¹	Percent Shading ¹		Number of Woody Debris		Percent Channel Flow ¹		Bank Stability ¹		Aesthetic Rating ¹	
Site		Epifaunal Substrate ¹		Pool Quality ¹		Percent Embeddedness ¹		Maximum Depth (cm) ¹		Number of Rootwads		Channel Alteration ¹		Riparian Width (m) ¹	
HA-P-087-207-97	17	18	17	17	14	40	50	91	3	1	95	9	11	50	14
HA-P-099-102-97	16	11	1	5	1	50	90	22	0	0	10	18	20	50	16
HA-P-100-204-97	15	14	18	18	16	40	80	77	1	0	95	15	15	0	15
HA-P-116-109-96	15	11	14	15	14	60	90	88	4	1	50	7	9	50	15
HA-P-128-104-96	6	7	8	7	8	75	90	29	0	0	75	6	6	5	11
HA-P-131-1-94	15	16	13	12	12	10	20	80	1		95	15	6	0	16
HA-P-131-3-94	6	4	11	13	11	60	60	55	1		95	2	10	0	15
HA-P-133-111-97	14	13	7	10	16	40	40	33	0	0	99	16	7	13	12
HA-P-138-302-97	16	15	13	15	5	95	45	93	5	3	99	13	8	50	10
HA-P-141-302-96	17	16	18	14	19	50	65	101	1	0	60	10	10	50	15
HA-P-141-303-96	14	12	15	15	14	40	40	77	3	0	98	15	10	22	14
HA-P-142-105-97	10	9	8	10	10	50	95	35	4	1	90	14	12	35	15
HA-P-151-102-96	8	11	6	6	10	65	98	47	0	0	95	1	11	0	0
HA-P-156-112-97	17	16	13	17	14	10	85	64	6	2	80	9	10	50	10
HA-P-164-306-96	12	5	13	13	13	80	60	75	1	10	90	6	5	0	11
HA-P-174-2-94	18	16	18	19	17	20	70	110	1		95	10	17	0	15
HA-P-174-4-94	17	17	18	15	19	20	50	104	0		95	15	19	30	17
HA-P-178-202-97	16	16	14	12	16	20	65	61	0	0	95	14	15	50	12
HA-P-178-209-97	18	18	18	16	18	15	90	91	4	3	99	18	18	50	17
HA-P-180-107-97	16	13	12	17	14	30	35	74	3	0	99	18	12	0	7
HA-P-186-308-97	18	10	18	16	5	40	60	102	3	3	75	6	10	50	15
HA-P-205-1-94	12	13	11	8	10	25	75	64	3		75	8	10	0	8
HA-P-205-2-94	18	15	13	14	14	30	95	68	12		70	8	7	10	14
HA-P-207-1-94	14	14	16	15	15	50	45	90	0		100	6	13	0	12
HA-P-207-2-94	17	17	16	13	15	40	60	73	2		85	8	12	0	15
HA-P-214-211-97	19	18	16	17	15	25	95	85	4	1	70	15	16	30	11
HA-P-216-1-94	3	4	1	1	1	50	95	10	1		50	3	4	0	2
HA-P-216-2-94	17	14	13	15	12	35	70	102	5		75	11	16	0	17
HA-P-225-1-94	12	11	9	7	11	50	90	43	4		55	7	6	50	15
HA-P-225-2-94	6	14	8	3	9	40	95	15	5		80	9	13	50	16
HA-P-244-1-94	12	11	8	8	10	50	65	38	2		85	8	11	1	16
HA-P-244-2-94	15	16	14	9	15	20	90	47	2		90	16	15	1	17
HA-P-246-304-96	12	11	16	14	14	55	60	85	5	6	85	11	11	50	16

¹ MBSS Qualitative Habitat Metric - See Appendix B for Guidance




Figure 5. Stream ecological conditions based on the Physical Habitat Index (PHI) at Maryland Biological Stream Survey sites in Harford County, 1994-1997.

Table 6. Fish Index of Biotic Integrity (F-IBI), Benthic Macroinvertebrate Index of Biotic Integrity (B-IBI), Family-Level Benthic Macroinvertebrate Index of Biotic Integrity (Fam. IBI), and Physical Habitat Index (PHI) scores at Maryland Biological Stream Survey sites in Harford County, 1994-1997.

Site	Stream Name	F-IBI	B-IBI	Fam. IBI	PHI
BA-P-042-116-96	Little Gunpowder Fls	4.11	3.7		86.97
BA-P-215-305-96	Little Gunpowder R	4.33	3.2		74.63
HA-N-003-204-96	Romney Cr Middle Br		1.3		
HA-N-009-105-96	Un Str		1.9		47.44
HA-N-018-103-96	Swan Cr		2.4		23.66
HA-N-036-206-96	Carsins Run	3.89	1.9		57.57
HA-N-040-307-96	Winters Run	3.44	2.7		78.34
HA-N-052-202-96	Gasheys Run	5.00	2.7		95.42
HA-N-067-111-96	Cranberry Run	3.22	1.0		91.84
HA-N-068-301-96	Bynum Run	4.56	2.4		96.97
HA-N-068-308-96	Bynum Run	4.56	1.0		71.55
HA-N-086-201-96	Broad Run	3.67	2.1		90.61
HA-N-092-304-96	Winters Run	4.11	3.0		87.02
HA-N-099-305-96	Bynum Run	4.56	1.6		93.48
HA-P-001-205-96	East Br	5.00	2.1		91.28
HA-P-005-206-97	Big Br	3.22	4.1		68.83
HA-P-008-1-94	Deep Cr	3.44	4.3		
HA-P-008-3-94	Deep Cr	4.33	4.1		
HA-P-010-103-97	South Stirrup Run	4.56	3.7		71.41
HA-P-012-205-97	Mine Br	4.56	3.0		89.12
HA-P-013-101-97	Herring Run	1.67	2.6		53.42
HA-P-024-116-97	Un Trib To Broad Cr		3.7		
HA-P-033-203-97	Little Deer Cr	5.00	3.2		93.44
HA-P-033-212-97	Little Deer Cr	4.56	3.9		57.46
HA-P-035-208-97	Plumtree Br	2.56	3.7		68.83
HA-P-037-104-97	Jacks Hole		3.4		57.46
HA-P-038-106-96	Heavenly Waters	4.56	1.4		62.38
HA-P-041-1-94	Deer Cr	1.89	3.4		
HA-P-041-2-94	Deer Cr			1.86	
HA-P-062-207-96	Bynum Run	2.56	1.9		5.25
HA-P-068-114-97	Big Br	2.78	3.4		69.27
HA-P-071-1-94	Deep Cr	4.56		3.57	
HA-P-071-2-94	Deep Cr	4.33		3.29	
HA-P-071-318-97	Broad Cr	4.56	3.9		89.12
HA-P-078-4-94	Cabbage Run Br	2.33		1.57	
HA-P-078-6-94	Cabbage Run Br	2.56		1.57	
HA-P-087-1-94	Stout Bottle Br	4.11		3.29	
HA-P-087-2-94	Stout Bottle Br	4.11		3.00	
HA-P-087-207-97	Cabbage Run Br	4.56	3.9		79.00
HA-P-099-102-97	Holland's Br		3.4		5.89
HA-P-100-204-97	Deer Cr	4.78	2.6		75.40
HA-P-116-109-96	Un Trib To Little Gunpowder R	3.22	3.2		62.38
HA-P-128-104-96	Un Trib To West Br Winters Run	2.33	3.4		5.78
HA-P-131-1-94	Deep Cr	4.56	2.3		
HA-P-131-3-94	Deep Cr	3.67	4.1		
HA-P-133-111-97	Un Trib To Deer Cr	2.78	3.7		38.29
HA-P-138-302-97	Broad Cr	4.56	3.0		37.81
HA-P-141-302-96	Winters Run	4.78	3.4		83.36
HA-P-141-303-96	Winters Run	4.78	3.2		58.46

Table 6 (cont.). Fish Index of Biotic Integrity (F-IBI), Benthic Macroinvertebrate Index of Biotic Integrity (B-IBI), Family-Level Benthic Macroinvertebrate Index of Biotic Integrity (Fam. IBI), and Physical Habitat Index (PHI) scores at Maryland Biological Stream Survey sites in Harford County, 1994-1997.

Site	Stream Name	F-IBI	B-IBI	Fam. IBI	PHI
HA-P-142-105-97	Un Trib To Broad Cr	1.67	3.4		22.89
HA-P-151-102-96	Plumtree Run	2.11	1.7		5.35
HA-P-156-112-97	Un Trib To Deer Cr		4.1		80.32
HA-P-164-306-96	Bynum Run	4.11	2.6		94.26
HA-P-174-1-94	Broad Cr			2.71	
HA-P-174-2-94	Broad Cr	4.78		2.14	
HA-P-174-4-94	Broad Cr				
HA-P-176-309-97	Broad Cr		4.6		
HA-P-178-202-97	Big Br	2.78	4.1		70.56
HA-P-178-209-97	Big Br	4.33	3.4		96.19
HA-P-180-107-97	Falling Br	3.44	3.9		51.38
HA-P-186-308-97	Broad Cr	5.00	3.4		77.61
HA-P-205-1-94	Broad Cr	3.22	1.2		
HA-P-205-2-94	Broad Cr	3.67	4.1		
HA-P-207-1-94	Stout Bottle Br	4.33		3.00	
HA-P-207-2-94	Stout Bottle Br	4.33		3.00	
HA-P-214-211-97	Holland Br	4.33	3.4		83.64
HA-P-216-1-94	Hopkins Br	1.22		1.57	
HA-P-216-2-94	Hopkins Br	5.00	3.4		
HA-P-225-1-94	Deep Cr	2.78	4.1		
HA-P-225-2-94	Deep Cr	1.44	3.7		
HA-P-244-1-94	Stout Bottle Br	3.44	3.7		
HA-P-244-2-94	Stout Bottle Br	3.67	2.6		
HA-P-246-304-96	Little Gunpowder R	4.11	3.7		91.12

Table 7. Water chemistry data collected at Maryland Biological Stream Survey sites in Harford County, 1994-1997.

Site	pH	Conductivity (μ S/cm)	Acid Neutralizing Capacity (μ eq/L)	Nitrate (mg/L)	Sulfate (mg/L)	Dissolved Oxygen (mg/L)	Dissolved Organic Carbon (mg/L)
BA-P-042-116-96	6.75	0.125	239.90	4.154	6.545	9.80	0.90
BA-P-215-305-96	7.78	0.140	515.60	2.687	7.153	11.20	2.30
HA-N-003-204-96	6.64	0.166	431.40	0.193	13.317		11.40
HA-N-009-105-96	6.81	1.433	669.80	1.096	21.332	8.80	5.20
HA-N-018-103-96	7.35	0.114	591.20	0.688	9.824	8.30	4.50
HA-N-036-206-96	9.04	0.182	510.80	1.018	9.435	9.30	3.40
HA-N-040-307-96	8.87	0.156	453.80	2.496	10.089	9.00	1.60
HA-N-052-202-96	7.37	0.151	429.00	2.318	11.055	11.00	2.10
HA-N-067-111-96	6.96	0.239	382.80	0.694	14.430	9.00	1.20
HA-N-068-301-96	7.66	0.232	807.10	2.246	17.904	10.00	2.40
HA-N-068-308-96	8.24	0.225	784.30	2.178	17.915	10.00	2.10
HA-N-086-201-96	7.95	0.142	450.60	3.254	11.122	11.90	2.30
HA-N-092-304-96	7.67	0.161	484.10	2.652	10.141	9.10	1.70
HA-N-099-305-96	8.35	0.202	666.30	2.248	15.964	10.30	2.00
HA-P-001-205-96	7.19	0.126	347.10	2.806	6.146	9.20	1.40
HA-P-005-206-97	7.04	0.119	223.10	5.996	4.694	10.10	0.50
HA-P-008-1-94	6.62	0.081	331.68	2.148	5.938		10.00
HA-P-008-3-94							
HA-P-010-103-97	7.11	0.099	322.60	2.112	10.934	9.60	1.60
HA-P-012-205-97	7.18	0.112	329.00	3.376	7.078	10.60	1.90
HA-P-013-101-97	7.21	0.199	389.80	1.586	13.662	9.50	2.60
HA-P-024-116-97	5.86	0.121	280.10	3.456	9.559		0.40
HA-P-033-203-97	7.24	0.124	415.50	3.402	9.221	9.60	1.30
HA-P-033-212-97	7.26	0.124	458.60	3.388	9.878	10.60	1.40
HA-P-035-208-97	7.29	0.152	319.50	3.408	6.567	9.70	1.20
HA-P-037-104-97	6.99	0.289	466.50	5.884	11.232	9.80	1.10
HA-P-038-106-96	7.93	0.276	1031.60	0.778	9.792	9.70	1.40
HA-P-041-1-94							
HA-P-041-2-94	6.43	0.126	335.08	2.333	12.509		1.00
HA-P-062-207-96	8.30	0.216	910.30	2.176	8.127	8.90	2.30
HA-P-068-114-97	6.94	0.122	258.10	5.944	5.635	9.80	1.00
HA-P-071-1-94	7.03	0.140	288.03	5.486	6.287		2.00
HA-P-071-2-94							
HA-P-071-318-97	7.15	0.180	313.00	4.772	8.719	11.10	1.70
HA-P-078-4-94	7.02	0.144	464.56	5.089	7.782		4.00
HA-P-078-6-94							
HA-P-087-1-94							
HA-P-087-2-94	7.49	0.138	400.38	4.843	9.367		3.00
HA-P-087-207-97	7.28	0.159	389.10	5.074	12.223	10.50	1.10
HA-P-099-102-97	7.10	0.089	395.00	0.588	11.304	7.70	3.40
HA-P-100-204-97	7.19	0.127	432.80	1.070	28.003	9.20	1.40
HA-P-116-109-96	7.43	0.145	488.20	2.063	12.124	9.80	2.40
HA-P-128-104-96	6.76	0.195	192.60	3.067	4.937	11.70	1.20
HA-P-131-1-94							
HA-P-131-3-94	6.77	0.082	178.39	3.476	4.753		2.00
HA-P-133-111-97	7.52	0.184	925.50	2.224	11.315	9.80	1.00
HA-P-138-302-97	7.29	0.144	382.10	3.608	8.035	9.40	1.30
HA-P-141-302-96	7.26	0.126	297.10	3.230	6.420	10.50	1.00

Table 7 (cont.). Water chemistry data collected at Maryland Biological Stream Survey sites in Harford County, 1994-1997.

Site	pH	Conductivity (μ S/cm)	Acid Neutralizing Capacity (μ eq/L)	Nitrate (mg/L)	Sulfate (mg/L)	Dissolved Oxygen (mg/L)	Dissolved Organic Carbon (mg/L)
HA-P-141-303-96	7.38	0.132	275.80	3.174	6.548	9.60	1.20
HA-P-142-105-97	6.91	0.150	340.50	4.738	10.286	9.30	0.80
HA-P-151-102-96	7.25	0.349	811.70	2.462	21.741	9.50	2.10
HA-P-156-112-97	6.84	0.125	279.40	3.252	5.286	8.90	1.70
HA-P-164-306-96	8.19	0.237	855.60	2.163	16.803	10.20	2.10
HA-P-174-1-94	7.43	0.133	269.67	4.776	5.255		2.00
HA-P-174-2-94							
HA-P-176-309-97	7.28	0.145	284.90	4.544	7.261		0.90
HA-P-178-202-97	7.10	0.119	258.90	4.588	6.062	10.00	1.40
HA-P-178-209-97	7.22	0.110	247.80	4.104	5.652	9.40	1.20
HA-P-180-107-97	7.04	0.141	314.60	4.766	6.016	9.60	1.90
HA-P-186-308-97	7.29	0.143	294.50	4.342	7.259	9.70	1.10
HA-P-205-1-94	7.18	0.139	634.37	1.573	9.993		2.00
HA-P-205-2-94							
HA-P-207-1-94	7.52	0.143	405.40	4.831	9.316		3.00
HA-P-207-2-94							
HA-P-214-211-97	7.47	0.143	671.60	2.338	11.443	10.50	2.30
HA-P-216-1-94							
HA-P-216-2-94	7.81	0.108	571.67	0.947	6.174		2.00
HA-P-225-1-94							
HA-P-225-2-94	6.49	0.070	306.49	1.808	4.721		2.00
HA-P-244-1-94	7.08	0.135	350.14	2.977	12.297		2.00
HA-P-244-2-94							
HA-P-246-304-96	7.56	0.137	456.80	2.596	6.958	9.70	1.50




Figure 6. Nitrate-nitrogen concentrations (mg/L) at Maryland Biological Stream Survey sites in Harford County, 1994-1997.

Literature Cited

- Barbour, M.T., and J.B. Stribling. 1991. Use of habitat assessment in evaluating the biological integrity of stream communities. In: *Biological Criteria: Research and Regulation*. U.S. Environmental Protection Agency, Washington, D.C. EPA-440/5-91-005. pp. 25-38.
- Cairns, J. and J.R. Pratt. 1993. A history of biological monitoring using benthic macroinvertebrates. In Rosenberg, D.M. and V.H. Resh, eds. *Freshwater Monitoring and Benthic Macroinvertebrates*. Chapman and Hall, New York
- CBRM (Chesapeake Bay Research and Monitoring Division). 1989. Acid deposition in Maryland. Summary of results through 1988. Maryland Department of Natural Resources, Chesapeake Bay Research and Monitoring Division, Annapolis, MD. AD-89-1. NTIS No. PB89-182729/AS.
- Chaillou, J.C. 1995. Technical Memorandum, The Maryland Biological Stream Survey: Summary of landowner contact procedures. Prepared by Versar, Inc., Columbia, MD, for Maryland Department of Natural Resources, Monitoring and Non-tidal Assessment Division, Annapolis, MD. CBWP-MANTA-EA-95-1.
- Cochran, W.G. 1977. *Sampling Techniques*. 3rd ed. New York: John Wiley and Sons.
- EPA (U.S. Environmental Protection Agency). 1987. *Handbook of methods for acid deposition studies: Laboratory analysis for surface water chemistry*. EPA-600/4-87/026.
- Hall, L.W., Jr., R.P. Morgan II, E.S. Perry, A. Waltz. 1999. Development of a provisional physical habitat index for Maryland freshwater streams. Prepared by Wye Research and Education Center, Queenstown, MD, for Maryland Department of Natural Resources, Monitoring and Non-tidal Assessment 455.
- Heimbuch, D.G., J.C. Seibel, H.T. Wilson, P.F. Kazyak. 1999. A multiyear lattice sampling design for Maryland-wide fish abundance estimation. *Journal of Agricultural, Biological, and Environmental Statistics*. 4: 443-455.
- Hilsenhoff, W.L. 1987. An improved biotic index of organic stream pollution. *Great Lakes Entomologist* 20: 31-39.
- Janicki, A., D. Wade, D. Heimbuch, H. Wilson, P. Jacobson, P. Kazyak. 1993. Maryland Biological Stream Survey design report. Prepared for Maryland Department of Natural Resources, Chesapeake Bay Research and Monitoring Division, by Coastal Environmental Services, Linthicum, MD. CBRM-AD-93-1.
- Jessen, R.J. 1978. *Statistical Survey Techniques*. John Wiley, New York.
- Karr, J.R. 1991. Biological integrity: A long-neglected aspect of water resource management. *Ecological Applications* 1: 66-84.
- Karr, J.R., K.D. Fausch, P.L. Angermeier, P.R. Yant, and I.J. Schlosser. 1986. Assessing biological integrity in running waters: a method and its rationale. *Illinois Natural History Survey Special Publication* 5. 28 pp.
- Kazyak, P.F. 1996. Maryland Biological Stream Survey Sampling Manual. Maryland Department of Natural Resources, Monitoring and Non-tidal Assessment Division.
- Kazyak, P.F. 1994. Maryland Biological Stream Survey Sampling Manual. Prepared by Versar, Inc., Columbia, MD, for Maryland Department of Natural Resources, Chesapeake Bay Research and Monitoring Division.
- Kerans, B.L. and J.R. Karr. 1994. A benthic index of biotic integrity (B-IBI) for rivers of the Tennessee valley. *Ecological Applications* 4: 768-785.
- Knapp, C.M., W.P. Saunders, D.G. Heimbuch, H.S. Greening, and G.J. Filbin. 1988. Maryland

- Synoptic Stream Chemistry Survey: Estimating the number and distribution of streams affected by or at risk from acidification. Prepared by International Science and Technology, Inc., Reston, VA, for the Maryland Department of Natural Resources, Power Plant Research Program, Annapolis, MD. AD-88-2. NTIS No. PB88-213996/AS.
- Knapp, C.M. and W.P. Saunders. 1987. Maryland Synoptic Stream Chemistry Survey design report. Prepared by International Science and Technology, Inc. for Maryland Department of Natural Resources, Power Plant Research Program. AD-87-5.
- Lenat, D.R. 1988. Water quality assessment of streams using a qualitative collection method for benthic macroinvertebrates. *Journal of North American Benthological Society* 7: 222-223.
- Miller, D.L., P.M. Leonard, R.M. Hughes, J.R. Karr, P.B. Moyle, L.H. Schrader, B.A. Thompson, R.A. Daniels, K.S. Fausch, G.A. Fitzhugh, J.R. Gammon, D.B. Halliwell, P.L. Angermeier, and D.J. Orth. 1988. Regional applications of an index of biotic integrity for use in water resource management. *Fisheries* 13(5): 12-20.
- Ohio EPA (Ohio Environmental Protection Agency). 1987. Biological criteria for the protection of aquatic life. Volumes I-III. Ohio EPA, Division of Water Quality Monitoring and Assessment, Surface Water Section, Columbus, Ohio.
- Plafkin, J.L., M.T. Barbour, K.D. Porter, S.K. Gross and R.M. Hughes. 1989. Rapid bioassessment protocols for use in streams and rivers: Benthic macroinvertebrates and fish. U.S. Environmental Protection Agency, Office of Water, Washington, D.C. EPA 440-4-89-001.
- Ranasinghe, J.A., S.B. Weisberg, D.M. Dauer, L.C. Schaffner, R.J. Diaz, and J.B. Frithsen. 1994. Chesapeake Bay Benthic Community Restoration Goals. Prepared by Versar, Inc. for U.S. EPA and the Maryland Department of Natural Resources, Chesapeake Bay Research and Monitoring Division.
- Rankin, E.T. 1989. The Qualitative Habitat Evaluation Index (QHEI): Rationale, methods, and application. Ohio EPA, Division of Water Quality Planning and Assessment, Ecological Analysis Section, Columbus, OH.
- Resh, V. H., 1995. Freshwater benthic macroinvertebrates and rapid assessment procedures for water quality monitoring in the developing and newly industrialized countries. In: Davis, W.S. and T.P. Simon, eds. 1995. Biological assessment and criteria: tools for water resource planning and decision making. Lewis Publishers, Boca Raton, FL.
- Roth, N.E., M.T. Southerland, G. Mercurio, J.C. Chaillou, P.F. Kazyak, S.A. Stranko, A.P. Prochaska, D.G. Heimbuch, and J.C. Seibel. 1999. State of the Streams: 1995-1997 Maryland Biological Stream Survey Results. Prepared by Versar, Inc., Columbia, MD, for Maryland Department of Natural Resources, Monitoring and Non-tidal Assessment Division. CBWP-MANTA-EA-99-6.
- Roth, N.E., M.T. Southerland, J.C. Chaillou, G.T. Wilson, D.G. Heimbuch, and J.C. Seibel. 1998a. Maryland Biological Stream Survey: Ecological Status of Non-tidal Streams Sampled in 1996. Prepared by Versar, Inc., Columbia, MD, and Post, Buckley, Schuh, and Jernigan, Inc., Bowie, MD, for Maryland Department of Natural Resources, Monitoring and Non-tidal Assessment Division. CBWP-MANTA-EA-97-2.
- Roth, N.E., M.T. Southerland, J.C. Chaillou, R.J. Klauda, P.F. Kazyak, S.A. Stranko, S.B. Weisberg, L.W. Hall, Jr., and R.P. Morgan II. 1998b. Maryland Biological Stream Survey: Development of a Fish Index of Biotic Integrity. *Environmental Management and Assessment* 51: 89-106.
- Roth, N.E., M.T. Southerland, J.C. Chaillou, J.H. Volstad, S.B. Weisberg, H.T. Wilson, D.G. Heimbuch, and J.C. Seibel. 1997. Maryland Biological Stream Survey: Ecological Status of Non-Tidal Streams in Six Basins Sampled in 1995. Prepared by Versar, Inc., Columbia, MD and Coastal Environmental Services, Linthicum, MD for Maryland Department of Natural Resources,

- Monitoring and Non-tidal Assessment Division.
CBWP-MANTA-EA-97-2.
- Simon, T.P. ed. 1999. Assessing the Sustainability and Biological Integrity of Water Resources Using Fish Communities. CRC Press, Washington DC.
- Strahler, A.N. 1957. Quantitative analysis of watershed geomorphology. Transactions of the American Geophysical Union 38(6): 913-920.
- Stribling, J.B., B.K. Jessup, J.S. White, D. Boward, M. Hurd. 1998. Development of a benthic index of biotic integrity for Maryland streams. Prepared by Tetra Tech, Inc., Owings Mills, MD, for Maryland Department of Natural Resources, Monitoring and Non-tidal Assessment Division, Annapolis, MD. CBWP-EA-98-3.
- Tornatore, T., P. Jacobson, J. Gerritsen and A. Janicki. 1992. Data sets for developing measures of biological integrity in Maryland streams. Prepared by Versar, Inc., Columbia, MD, for the Maryland Department of Natural Resources, Chesapeake Bay Research and Monitoring Division, Annapolis, MD. AD-93-7.
- Weisberg, S.B., J.A. Ranasinghe, D.M. Dauer, L.C. Schaffner, R.J. Diaz, J.B. Frithsen. 1997. An estuarine benthic index of biotic integrity (B-IBI) for Chesapeake Bay. Estuaries 20: 149-158.

**THIS PAGE INTENTIONALLY
LEFT BLANK**

Appendix A. Summary of the types of data collected at Maryland Biological Stream Survey sites in Harford County, 1994-1997. Abbreviations used are as follows: F-IBI - Fish Index of Biotic Integrity; B-IBI Benthic Macroinvertebrate Index of Biotic Integrity; Fam.IBI - Family-Level Benthic Macroinvertebrate Index of Biotic Integrity; PHI - Physical Habitat Index.

Site	Stream Name	Benthic Macroinvertebrate		Habitat		F-IBI		Fam. IBI	
		Fish	Herpetofauna	Water Chemistry			B-IBI	PHI	
BA-P-042-116-96	Little Gunpowder Fls	X	X	X	X	X	X	X	X
BA-P-215-305-96	Little Gunpowder R	X	X	X	X	X	X	X	X
HA-N-003-204-96	Romney Cr Middle Br		X		X		X		
HA-N-009-105-96	Un Str	X	X	X	X		X		X
HA-N-018-103-96	Swan Cr	X	X	X	X		X		X
HA-N-036-206-96	Carsins Run	X	X	X	X	X	X		X
HA-N-040-307-96	Winters Run	X	X	X	X	X	X		X
HA-N-052-202-96	Gasheys Run	X	X	X	X	X	X		X
HA-N-067-111-96	Cranberry Run	X	X	X	X	X	X		X
HA-N-068-301-96	Bynum Run	X	X	X	X	X	X		X
HA-N-068-308-96	Bynum Run	X	X	X	X	X	X		X
HA-N-086-201-96	Broad Run	X	X	X	X	X	X		X
HA-N-092-304-96	Winters Run	X	X	X	X	X	X		X
HA-N-099-305-96	Bynum Run	X	X	X	X	X	X		X
HA-P-001-205-96	East Br	X	X	X	X	X	X		X
HA-P-005-206-97	Big Br	X	X	X	X	X	X		X
HA-P-008-1-94	Deep Cr	X	X	X	X	X	X		
HA-P-008-3-94	Deep Cr	X	X	X	X	X	X		
HA-P-010-103-97	South Stirrup Run	X	X	X	X	X	X		X
HA-P-012-205-97	Mine Br	X	X	X	X	X	X		X
HA-P-013-101-97	Herring Run	X	X	X	X	X	X		X
HA-P-024-116-97	Un Trib To Broad Cr		X		X		X		
HA-P-033-203-97	Little Deer Cr	X	X	X	X	X	X		X
HA-P-033-212-97	Little Deer Cr	X	X	X	X	X	X		X
HA-P-035-208-97	Plumtree Br	X	X	X	X	X	X		X
HA-P-037-104-97	Jacks Hole	X	X	X	X		X		X
HA-P-038-106-96	Heavenly Waters	X	X	X	X	X	X		X
HA-P-041-1-94	Deer Cr	X	X	X	X	X	X		
HA-P-041-2-94	Deer Cr	X	X	X	X			X	
HA-P-062-207-96	Bynum Run	X	X	X	X	X	X		X
HA-P-068-114-97	Big Br	X	X	X	X	X	X		X
HA-P-071-1-94	Deep Cr	X	X	X	X			X	
HA-P-071-2-94	Deep Cr	X	X	X		X		X	
HA-P-071-318-97	Broad Cr	X	X	X	X	X	X		X
HA-P-078-4-94	Cabbage Run Br	X	X	X	X	X		X	

Appendix A (cont.). Summary of the types of data collected at Maryland Biological Stream Survey sites in Harford County, 1994-1997. Abbreviations used are as follows: F-IBI - Fish Index of Biotic Integrity; B-IBI - Benthic Macroinvertebrate Index of Biotic Integrity; Fam. IBI - Family-Level Benthic Macroinvertebrate Index of Biotic Integrity; PHI - Physical Habitat Index.

Site	Stream Name	Benthic Macroinvertebrate		Habitat		F-IBI		Fam. IBI	
		Fish	Herpetofauna	Water Chemistry			B-IBI		PHI
HA-P-078-6-94	Cabbage Run Br	X	X	X	X		X	X	
HA-P-087-1-94	Stout Bottle Br	X	X	X	X		X	X	
HA-P-087-2-94	Stout Bottle Br	X	X	X	X	X	X	X	
HA-P-087-207-97	Cabbage Run Br	X	X	X	X	X	X		X
HA-P-099-102-97	Holland's Br	X	X	X	X	X	X		X
HA-P-100-204-97	Deer Cr	X	X	X	X	X	X		X
HA-P-116-109-96	Un Trib To Little Gunpowder R	X	X	X	X	X	X		X
HA-P-128-104-96	Un Trib To West Br Winters Run	X	X	X	X	X	X		X
HA-P-131-1-94	Deep Cr	X	X	X	X	X	X		
HA-P-131-3-94	Deep Cr	X	X	X	X	X	X		
HA-P-133-111-97	Un Trib To Deer Cr	X	X	X	X	X	X		X
HA-P-138-302-97	Broad Cr	X	X	X	X	X	X		X
HA-P-141-302-96	Winters Run	X	X	X	X	X	X		X
HA-P-141-303-96	Winters Run	X	X	X	X	X	X		X
HA-P-142-105-97	Un Trib To Broad Cr	X	X	X	X	X	X		X
HA-P-151-102-96	Plumtree Run	X	X	X	X	X	X		X
HA-P-156-112-97	Un Trib To Deer Cr	X	X	X	X		X		X
HA-P-164-306-96	Bynum Run	X	X	X	X	X	X		X
HA-P-174-1-94	Broad Cr		X		X			X	
HA-P-174-2-94	Broad Cr	X	X	X	X	X		X	
HA-P-174-4-94	Broad Cr	X		X	X				
HA-P-176-309-97	Broad Cr		X		X		X		
HA-P-178-202-97	Big Br	X	X	X	X	X	X		X
HA-P-178-209-97	Big Br	X	X	X	X	X	X		X
HA-P-180-107-97	Falling Br	X	X	X	X	X	X		X
HA-P-186-308-97	Broad Cr	X	X	X	X	X	X		X
HA-P-205-1-94	Broad Cr	X	X	X	X	X	X		
HA-P-205-2-94	Broad Cr	X	X	X	X	X	X		
HA-P-207-1-94	Stout Bottle Br	X	X	X	X	X		X	
HA-P-207-2-94	Stout Bottle Br	X	X	X	X	X		X	
HA-P-214-211-97	Holland Br	X	X	X	X	X	X		X
HA-P-216-1-94	Hopkins Br	X	X	X	X	X		X	
HA-P-216-2-94	Hopkins Br	X	X	X	X	X	X		
HA-P-225-1-94	Deep Cr	X	X	X	X	X	X		
HA-P-225-2-94	Deep Cr	X	X	X	X	X	X		

Appendix A (cont.) Summary of the types of data collected at Maryland Biological Stream Survey sites in Harford County, 1994-1997.
 Abbreviations used are as follows: F-IBI - Fish Index of Biotic Integrity; B-IBI - Benthic Macroinvertebrate Index of Biotic Integrity; Fam. IBI - Family-Level Benthic Macroinvertebrate Index of Biotic Integrity; PHI - Physical Habitat Index.

Site	Stream Name	Benthic Macroinvertebrate			Habitat	F-IBI		Fam. IBI	PHI
		Fish	Herpetofauna			Water Chemistry	B-IBI		
HA-P-244-1-94	Stout Bottle Br	X	X	X	X	X	X	X	
HA-P-244-2-94	Stout Bottle Br	X	X	X	X		X	X	
HA-P-246-304-96	Little Gunpowder R	X	X	X	X	X	X	X	X

**THIS PAGE INTENTIONALLY
LEFT BLANK**

Appendix B. Physical habitat condition measured by the Maryland Biological Stream Survey, 1994-1997. All variables rated on a scale of 0 (poor) to 20 (optimal) unless otherwise noted.

SUBSTRATE AND INSTREAM COVER

Instream Habitat is rated according to the perceived value of habitat to the fish community. Higher scores are assigned to sites with a variety of habitat types and particle sizes. In addition, higher scores are assigned to sites with a high degree of uneven substrate, including logs and rootwads. In streams where substrate types are favorable but flows are so low that fish are essentially precluded from using the habitat, low scores are assigned. If none of the habitat within a segment is useable by fish, a score of zero is assigned.

Epifaunal Substrate is rated based on the amount and variety of hard, stable substrates usable by benthic macroinvertebrates. Because they inhibit colonization, flocculent materials or fine sediments surrounding otherwise good substrates are assigned low scores. Scores are also reduced when substrates are less stable.

Velocity/Depth Diversity is rated based on the variety of velocity/depth regimes present at a site (slow-shallow, slow-deep, fast-shallow, and fast-deep). As with embeddedness, this metric varies by stream gradient.

Pool/Glide/Eddy Quality is rated based on the variety and spatial complexity of slow or still water habitat within the sample segment. In high-gradient streams, functionally important slow water habitat may exist in the form of larger eddies. Within a category, higher scores are assigned to segments which have undercut banks, woody debris or other types of cover for fish.

Riffle/Run Quality is based on the depth, complexity, and functional importance of riffle/run habitat in the segment, with highest scores assigned to segments dominated by deeper riffle/run areas, stable substrates, and a variety of current velocities.

Embeddedness is a percentage of surface area of larger particles that is surrounded by fine sediments on the stream bottom. In low gradient streams, embeddedness may be high even in relatively unimpaired watersheds.

CHANNEL CHARACTER

Channel Alteration is a measure of large-scale changes in the shape of the stream channel. Channel alteration includes: concrete channels, artificial embankments, obvious straightening of the natural channel, rip-rap, or other structures, as well as recent bar development. Ratings for this metric are based on the presence of artificial structures as well as the existence, extent, and coarseness of point bars, side bars, and mid-channel bars which indicate the degree of flow fluctuations and substrate stability. Evidence of channelization may sometimes be seen in the form of berms that parallel the stream channel.

Bank Stability is rated based on the presence/absence of riparian vegetation and other stabilizing bank materials such as boulders and rootwads, and frequency/size of erosional areas. Sites with steep slopes are not penalized if banks are composed solely of stable materials.

Channel Flow Status is the percentage of the stream channel that has water, with subtractions made for exposed substrates and dewatered areas.

RIPARIAN CORRIDOR

Shading is rated based on estimates of the degree and duration of shading at a site during summer, including any effects of shading caused by land forms.

Appendix B (cont.).Physical habitat condition measured by the Maryland Biological Stream Survey, 1994-1997.
All variables rated on a scale of 0 (poor) to 20 (optimal) unless otherwise noted.

Riparian Buffer is rated according to the size and type of the vegetated riparian buffer zone at the site. Cultivated fields for agriculture that have bare soil to any extent are not considered as riparian buffers. At sites where the buffer width is variable, or direct delivery of storm runoff or sediment to the stream is evident or highly likely, the narrowest representative buffer width in the segment (e.g., 0 if parking lot runoff enters directly to the stream) is measured and recorded even though some of the stream segment may have a well developed riparian buffer.

AESTHETICS/REMOTENESS

Aesthetics are rated according to the visual appeal of the site and presence/absence of human refuse, with highest scores assigned to stream segments with no human refuse and visually outstanding character.

Remoteness is rated based on the absence of detectable human activity and difficulty in accessing the segment.